

D R O P

**DIGITALE. 142.** Shell conical, sub-turreted, glabrous, with the aperture effuse at the base.

*Buccinum digitale.* *Meuschen Mus. Gevers.* p. 296. No. 507.

*Buccinum vittatum*, Var. *Gmelin*, p. 3500. *Schreibers Conch. i.* p. 168.

*Buccinum*, No. 180. *Schroeter Einl. i.* p. 404.

*Lister Conch.* t. 977. f. 33. *Petiver Gaz.* t. 102. f. 15.

*Adanson Senegal*, t. 4. f. 1. *Martini*, iv. p. 310. t. 155. f. 1468 and 1469, and t. 157. f. 1491.

*Variety.* Narrower, and somewhat elongated.

*Buccinum*, No. 181, and No. 182. *Schroeter Einl. i.* p. 405.

*Adanson Senegal*, t. 4. f. 2. *Knorr*, v. t. 22. f. 5. *Martini*, iv. t. 155. t. 1470 and 1471. *Favanne*, t. 40. f. C 1.

Inhabits the coasts of Bombay. *Petiver.* *Senegal.* *Adanson.* *Coromandel.* *Martini.*

Shell about an inch and a half long, and more than one-third as broad, whitish or brown, with a paler or bluish band at the upper margin of the whorls. I have followed Kæmmerer, Schroeter, and Martini, in arranging this as a separate species, and it differs from *B. vittatum* in not having either any transverse striæ or crenatures at the upper margin of the whorls. *Martini*, f. 1491, probably differs only in being coated with a green epidermis.

DILLWYN 1817 h 46

MIRAN

RAFFEL

NOT IN BRITISH MUSEUM

NEVER ASSIGNED TO THIS GENUS

NEVER ASSIGNED TO THE GENUS TERRERA