

THE
ZOOLOGY
OF THE
VOYAGE OF H.M.S. SAMARANG;

UNDER THE COMMAND OF
CAPTAIN SIR EDWARD BELCHER, C.B., F.R.A.S., F.G.S.,

DURING THE YEARS 1843-1846.

BY
JOHN EDWARD GRAY, F.R.S.; SIR JOHN RICHARDSON, M.D., F.R.S.;
ARTHUR ADAMS, F.L.S.; LOVELL REEVE, F.L.S.;
AND
ADAM WHITE, F.L.S.

EDITED BY ARTHUR ADAMS, F.L.S.,
ASSISTANT-SURGEON TO THE EXPEDITION.



Published under the Authority of the Lords Commissioners of the Admiralty.

LONDON:
REEVE AND BENHAM, 5, HENRIETTA STREET, COVENT GARDEN.
1850.

8. TERE BRA.

1. *TEREBRA SEROTINA*. Pl. X. Fig. 20. Ter. testâ lanceolato-turritâ, basi breviter recurvâ, anfractibus planulatis, supernè plicato-nodulosis, deinde arcuatim liratis, liris striis transversis numerosis irregularibus decussatis; citrino-aurantiâ.

HAB. Japan Island, Nangasaki Bay.

The noduled sculpture round the edge of the whorls, forming somewhat of a shelf, makes the sutures very distinct. The nodules are slightly plicate and pass into arched concentric ridges.

2. *TEREBRA ALBICOSTATA*. Pl. X. Fig. 21. Ter. testâ subulatâ, anfractibus supernè plicato-nodosis, deinde costatis, costis angustis, subirregularibus, interstitiis transversim impresso-striatis; castaneo-rufâ, costis nodisq̄ue albidis, anfractu ultimo basin versus albifasciato.

HAB. China Sea.

The ribs and nodules, although naturally white, appear colourless from the effect of abrasion.

3. *TEREBRA CÆLATA*. Pl. X. Fig. 22. Ter. testâ lanceolato-turritâ, anfractibus longitudinaliter arcuatim plicato-costatis, costis liris duplicatis striisque transversis cancellatis, sulco unico conspicuo infra suturas; albâ.

HAB. Philippine Islands.

The chief peculiarity of this delicately carved species consists in the transverse ridges being finely duplicate.

4. *TEREBRA AREOLATA*. Pl. X. Fig. 23. Ter. testâ lanceolato-turritâ, gracili, anfractibus planis, longitudinaliter plicato-costatis, costis tumidiusculis, confertis, subundulatis, interstitiis alveolatis; rubellâ, albo variegatâ.

HAB. China Sea.

The opaque white marbling of the shell is mainly upon the ribs, which have a swollen appearance, and are unusually close-set.

5. *TEREBRA ROSEATA*. Pl. X. Fig. 24. Ter. testâ subulatâ, anfractibus plicato-costatis, costarum interstitiis liris tenuibus clathratis, sulco prominulo infra suturas; pallidè roseâ.

HAB. Sooloo Islands; from sandy mud, at a depth of about thirty fathoms.

Of a uniform, delicate, transparent, pink colour.

6. *TEREBRA TORQUATA*. Pl. X. Fig. 13. Ter. testâ lanceolato-turritâ, anfractibus concavis, arcuatim costatis, supernè biseriatis, infra uniseriatis, nodosis, transversim striatis; fuscescente-albâ, ferrugineo marmoratâ.

HAB. China Sea.

