

SMITHSONIAN INSTITUTION.  
UNITED STATES NATIONAL MUSEUM.

---

# PROCEEDINGS

OF THE

UNITED STATES NATIONAL MUSEUM.

---

Volume XXIV.

---

PUBLISHED UNDER THE DIRECTION OF THE SMITHSONIAN INSTITUTION.



WASHINGTON:  
GOVERNMENT PRINTING OFFICE.  
1902.

East Florida, Bermuda, south to Cuba and west to Yucatan. U.S.N.M., 94431.

The figured specimen is from Bermuda, collected by Goode, and is what Jones referred to *S. brunnea* Hanley. It appears to be merely a better-nourished, slightly darker-colored geographical race of the *alternata* of Say. If the *S. picta* Orbigny (1842) comprises only a single species, which is not yet certain, this species extends southward to Rio Janeiro, Brazil. It is somewhat singular that no good figure of Say's shell is to be found in the whole conchological literature of America, though the shell has been known (chiefly by bleached, worn specimens) for three-quarters of a century.

EAST AMERICAN MARINE SPECIES.

CYLICHNA VERRILLII Dall.

Plate XXIX, fig. 1.

*Cylichna verrillii* DALL, Rep. Blake Gastr., Bull. Mus. Comp. Zool., XVIII, Jan., 1889, p. 54.

Collected off the coast of North Carolina at various stations in from 50 to 124 fathoms, sand, with the bottom temperatures from 58° to 75° F., by the U. S. Fish Commission steamer *Albatross*. Bluish white with a nearly translucent periostracum and finely spirally striated. U.S.N.M., 94477.

RETUSA MAYOI Dall.

Plate XXIX, fig. 2.

*Utriculus mayoi* DALL, Rep. Blake Gastr., Bull. Mus. Comp. Zool., XVIII, Jan., 1889, p. 46.

Collected from the stomach of a haddock at Portland, Maine, by Mr. John Mayo. U.S.N.M., 95269.

Shell solid, yellowish white, with little or no spiral striation.

UMBRACULUM (HYALOPATINA) RUSHII Dall.

Plate XXX, fig. 5.

*Umbraculum (Hyalopatina) rushii* DALL, Rep. Blake Gastr., Bull. Mus. Comp. Zool., XVIII, 1889, p. 61.—PILSBRY, Man. Conch., XVI, 1896, p. 185, pl. LI, fig. 58.

Dredged off Great Isaac Light, Bahamas, in 30 fathoms, sand, by Dr. W. H. Rush, U. S. N. U.S.N.M., 61222.

The shell is almost perfectly flat and very thin; bluish translucent white, dextral with a sinistral nucleus.

TEREBRA TEXANA Dall.

Plate XXIX, fig. 8.

*Terebra texana* DALL, Nautilus, XII, No. 4, Aug., 1898, p. 45.

From the Gulf coast of Matagorda Island, Texas, J. D. Mitchell. U.S.N.M., 107373.

Color pale yellowish,  
Only two imperfect speci-  
from the above locality.

TEREBRA

*Subula floridana* DALL, Rep.  
p. 63.

*Terebra (Subula) floridana* DALL

Key West and Florida  
sion steamer *Albatross* in  
It is of a pale straw color.

TEREBRA

*Terebra (Acus) rushii* DALL  
1889, p. 64.

Dredged 5 miles off Cay  
U. S. N. U.S.N.M., 829  
It is a small, brilliant  
spiral sculpture.

CONUS

*Conus stimpsoni* DALL, MS.

Dredged by the U. S.  
fathoms, off Key West.

Shell of about twelve  
what concave, turreted, t  
lowing four furnished v  
becomes entire on the s  
whorls are slightly conca  
three faint spiral groov  
shallow squarish channel  
straw color, finely axially  
white, suffused with salmon  
and base, with a very faint  
between base and should

The species is named  
whose last field work wa

DAPHNEL

Dredged at station 232  
ature 78° F., by the U.

Color pale yellowish, with darker yellow or brown flammules. Only two imperfect specimens of this fine species are known, both from the above locality.

TEREBRA (SUBULA) FLORIDANA Dall.

Plate XXIX, fig. 9.

*Subula floridana* DALL, Rep. Blake Gastr., Bull. Mus. Comp. Zool., XVIII, 1889, p. 63.

*Terebra (Subula) floridana* DALL, Bull. 37, U. S. Nat. Mus., 1889, p. 94.

Key West and Florida Strait, dredged by the U. S. Fish Commission steamer *Albatross* in 45 to 56 fathoms, sand. U.S.N.M., 87222. It is of a pale straw color.

TEREBRA (ACUS) RUSHII Dall.

Plate XXIX, fig. 6.

*Terebra (Acus) rushii* DALL, Rep. Blake Gastr., Bull. Mus. Comp. Zool., XVIII, 1889, p. 64.

Dredged 5 miles off Cape Florida, in 8 fathoms, by Dr. W. H. Rush, U. S. N. U.S.N.M., 82952.

It is a small, brilliantly white shell, remarkable for its exclusively spiral sculpture.

CONUS STIMPSONI, new species.

Plate XXIX, fig. 7.

*Conus stimpsoni* DALL, MS., in Coll. U. S. Nat. Mus.

Dredged by the U. S. Fish Commission steamer *Albatross*, in 60 fathoms, off Key West. U.S.N.M., 107371.


Shell of about twelve whorls, as figured, slopes of the spire somewhat concave, turreted, the nuclear whorl rounded, smooth; the following four furnished with a beaded keel at the shoulder; this keel becomes entire on the subsequent whorls; above the shoulder the whorls are slightly concave, the suture appressed, and there are about three faint spiral grooves on the concave surface; last whorl with shallow squarish channels, as figured. The periostracum is thin, pale straw color, finely axially striated. The color of the shell is pinkish white, suffused with salmon pink near the shoulder and on the spire and base, with a very faint, cloudy band of the same about midway between base and shoulder. Altitude of shell, 37 mm.

The species is named in honor of the late Dr. William Stimpson, whose last field work was done in the vicinity of the Florida Strait.

DAPHNELLA EUGRAMMATA, new species.

Plate XXIX, fig. 3.

Dredged at station 2328, off Habana, in 203 fathoms, bottom temperature 78° F., by the U. S. Fish Commission. U.S.N.M., 107448.



EAST AMERICAN GASTROPODS.

FOR EXPLANATION OF PLATE SEE PAGE 563.