


BULLETIN OF THE INSTITUTE OF MALACOLOGY TOKYO



83

INSTITUTE OF MALACOLOGY OF TOKYO
6-36, MIDORICHO 3 CHOME,
TANASHI CITY, TOKYO 188, JAPAN

VOLUME 2, NUMBER 2
April 30, 1985

Bull. Inst. Malac. Tokyo.
2(2): 24-27 (1985)

Newly Recorded or Noteworthy Molluscs
from Okinawa Islands (Ryukyu Group) (1)
Sadao Kosuge
(Institute of Malacology of Tokyo)
(Plate 11)

Angaria vicdani Kosuge, 1980 (pl. 11, f. 1)

1980 *Angaria vicdani* Kosuge, Bull. Inst. Malac. Tokyo, 1(3): 38-39, pl. 7, figs. 4-7; pl. 8,
figs. 4-8.

1982 *Angaria vicdani*: Abbott and Dance, Comp. Seashells, p. 51.

Hitherto known localities: Davao, Mindanao; Bohol and its adjacent area.

The present material was collected from Tibishi Point, off Okinawa Main Island at
about 35 m deep.

Measurements of the present material:

	Height(mm)	Breadth(mm)
No. 1	31.0	49.8
No. 2	31.0	50.4
No. 3	36.0	57.1
No. 4	29.0	49.3

Cypraea martini Schepman, 1907 (pl. 11, figs. 2,7)

1907 *Cypraea martini* Schepman, Samml. Geol. Reichs-Mus., Leiden, 1(8): 184, pl. 11, f. 4.

1930 *Adusta martini superstes* Schilder, Zool. Anz., 87: 114.

1932 *Erronea (Adusta) martini martini*: Schilder, Foss. Cat., 55: 187.

1952 *Notadusta martini*: Schilder and Schilder, Mem. Inst. Roy. Nat. Hist. Belg., 2(45):
137.

1963 *Cypraea (Notadusta) martini*: Cate, Veliger, 6: 80-84, pl. 5, figs. 1-4.

1966 *Notadusta martini martini*: Cate, ibid, 8: 200, pl. 29, figs. 5-6.

1967 *Cypraea martini superstes*: Cernohorsky, Mar. Shells Pacific, p. 91, pl. 16, f. 85.

1970 *Cypraea martini*: Burgess, Liv. Cow., pp. 60-61, pl. 4, f. D.

1980 *Cypraea martini*: Abbott and Dance, Comp. Seashells, p. 90.

Hitherto known localities: N. W. Celebes (Pleistocene fossil); Banuatsu; New Caledonia;
Philippines; Queensland.

Measurements of the present material:

Shell height (mm): 17.0; 16.7; 16.2; 16.2; 16.4; 15.8; 15.5; 15.3.

The present material was collected off Manzamo Cape, Okinawa Main Island at about
30-35 m deep by the scuba diving. There are fairly good amount of the specimens, but it
is very hard to get alive one.

Marchia laqueata (Sowerby, 1841) (pl. 11, f. 4)

1841 *Murex laqueatus* Sowerby, Conch. Illust., p. 3, sp. 45, pl. 190, f. 78.

1879 *Murex laqueatus*: Sowerby, Thes. Conchyl., pl. 9, f. 78.

1976 *Marchia laqueatus*: Radwin and D'Attilio, Murex Shell World, pp. 58-59, pl. 9, f. 5.

1982 *Conus legatus*: Abbott and Dance, Comp. Seashells, p. 245.
1984 *Conus legatus*: Kosuge, Bull. Inst. Malac. Tokyo, 1(10): 151, pl. 53, f. 14.
Hitherto known localities: New Caledonia~Philippines~Kii Peninsula, Japan.
The present material, measured 29.6 × 12.3 mm, was collected from Chibishi Point,
Okinawa Main Island at about 20 m deep.

Terebra fujitai Kuroda and Habe, 1952 (pl. 11, f. 11)

1952 *Terebra fujitai* Kuroda and Habe, Check List Bibliog. Rec. Mar. Moll. Jap., p. 89.
1960 *Terebra (Myurella) fujitai*: Azuma, Cat. Shell-bear. Moll. Tosa, p. 57, sp. 1067.
1963 *Terebra (Myurella) fujitai*: Shikama and Horikoshi, Select. Shells World, 1: 126, pl.
102, f. 9.
1965 *Terebra (Myurella) fujitai*: Nakayama, Cat. Moll. Tosa (Kochi), p. 63, sp. 1491.
1979 *Myurella fujitai*: Lan, Rare Shell Taiwan Col., p. 83, sp. 79.
1979 *Myurella fujitai*: Matsumoto, Moll. Shells Mie Pref., p. 69, sp. 1203.
Hitherto known localities: Sagami Bay~Kii Peninsula~Tosa Bay, Japan; East China
Sea.
The present material, measured 128.1 × 15.9 mm, was collected at Chibishi Point,
Okinawa Main Island at about 35 m deep.

Emendations of Two Specific Names

Sadao Kosuge

(Institute of Malacology of Tokyo)

The following species proposed in this series had typographical errors:

- Pterynotus cerinamarumai* Kosuge, 1980
Bull. Inst. Malac. Tokyo 1(4): 53-54, pl. 14, figs. 3, 5-9; pl. 15, figs. 1-2.
Latiaxis cerinamarumai Kosuge, 1980
Ibid., 1(4): 59-60, pl. 15, figs. 6-7; pl. 16, figs. 1-6.
The author emends both of them as follows:
Pterynotus celinamarumai nom. nov.
Latiaxis celinamarumai nom. nov.

Explanation of Plates

Plate 9

Figs. 1-10. *Lunatica choshiensis* Tiba, n. sp. (see p. 19): figs. 1, 3 off Kitami, Hokkaido (48 × 42 mm); f. 2 off Kuji, Iwate Pref. (15 × 13 mm); f. 4 off kuji, Iwate Pref. (250 m deep) (18 × 15 mm); f. 5 Paratype (46 × 40 mm); figs. 6-10 off Erimo, Hokkaido (200-250 m deep); f. 6 (20 × 17 mm); f. 7 (19 × 16 mm); f. 8 Holotype (30 × 24 mm); f. 9 (20 × 17 mm); f. 10 (19 × 17 mm).

Plate 10

Figs. 1-8. *Tectonatica rikuzenensis* Tiba, n. sp. (see p. 20): figs. 1-5 off Rikuzen, Iwate Pref., figs. 1-2, 4 Paratype (18 × 17 mm); figs. 3, 5 Holotype (14 × 13 mm); figs. 6-8 off Choshi, Tiba Pref., figs. 6, 7 (15 × 15 mm); f. 8 (11 × 9 mm).

Plate 11

Figs. 1-13 Molluscs from Okinawa Islands (see p. 24): f. 1 *Angaria viedani* (31.0 × 49.8 mm); figs. 2, 7 *Cypraea martini* (17.0 mm); f. 3 *Marchia bibbeyi* (20.3 × 14.2 mm); f. 4 *Marchia laqueata* (28.9 × 17.7 mm); f. 5 *Siratus alabaster* (133.8 × 88.3 mm); f. 6 *Conus tenuistriatus* (23.5 × 8.6 mm); f. 8 *Conus otohimeae* (20.7 × 10.0 mm); f. 9 *Conus legatus* (29.6 × 12.3 mm); f. 10 *Harpa kajiyamai* (51.4 mm); f. 11 *Terebra fujitai* (128.1 × 15.9 mm); f. 12 *Conus kintoki* (95.0 × 44.7 mm); f. 13 *Conus teramachii* (106.2 × 46.7 mm).

Plate 12

Figs. 1-4 New Species of the genus *Bolma* from Philippines (see p. 23): figs. 1-2 Holotype (10.1 × 11.7 mm); figs. 3-4 Paratype (9.4 × 11.3 mm). Figs. 5-10 Cypraeidae from Philippines (see p. 28): f. 5 *Cyp. guttata* (65.7 × 40.3 mm); f. 6 *Cyp. teramachi* (59.4 × 35.5 mm); f. 7 *Cyp. kuroharai* (47.1 × 33.3 mm); f. 8 *Cyp. porteri* (57.2 × 38.9 mm); f. 9 *Cyp. leucodon* (81.3 × 55.8 mm); f. 10 *Cyp. valentia* (92.9 × 68.3 mm).

Plate 13

Figs. 1-19 Cypraeidae from Philippines (see p. 28): figs. 1-2 *Cyp. ventriculus*; f. 3 *Cyp. onyx*; figs. 4-5 *Cyp. maculifera*; figs. 6-7 *Cyp. vitellus*; figs. 8-10 *Cyp. tigris*; figs. 11, 13 *Cyp. testudinaria*; f. 12 *Cyp. aurantium*; f. 14 *Cyp. leviathan*; f. 15 *Cyp. talpa*; figs. 16-17 *Cyp. mauritiana*; f. 18 *Cyp. argus*; f. 19 *Cyp. lynx*.

Plate 14

Figs. 1-36 Cypraeidae from Philippines (see p. 28): f. 1 *Cyp. errones*; f. 2 *Cyp. ovum*; figs. 3-4 *Cyp. walkeri*; f. 5 *Cyp. cylindrica*; f. 6 *Cyp. carneola*; f. 7 *Cyp. isabella*; f. 8 *Cyp. teres*; f. 9 *Cyp. stolida*; f. 10 *Cyp. helvola*; f. 11 *Cyp. poraria*; f. 12 *Cyp. labrolineata*; figs. 13-14 *Cyp. limacina*; f. 15 *Cyp. staphylaea*; f. 16 *Cyp. pyriformis*; figs. 17-18 *Cyp. pulchella*; figs. 19-20 *Cyp. moneta*; f. 21 *Cyp. annulus*; f. 22 *Cyp. cibraria*; f. 23 *Cyp. boivini*; f. 24 *Cyp. cuatoni*; f. 25 *Cyp. caputserpentis*; f. 26 *Cyp. depressa*; figs. 27-28 *Cyp. erosa*; f. 29 *Cyp. quadrimaculata*; f. 30 *Cyp. scurra*; f. 31 *Cyp. miliaris*; f. 32 *Cyp. arabica*; figs. 33-34 *Cyp. eglantina*; f. 35 *Cyp. caurica*; f. 36 *Cyp. chinensis*.

Plate 15

Figs. 1-34 Cypraeidae from Philippines (see p. 28): f. 1 *Cyp. ziczac*; f. 2 *Cyp. lutea*; figs. 3-4 *Cyp. nucleus*; f. 5 *Cyp. childreni*; f. 6 *Cyp. globulus*; f. 7 *Cyp. cicercula*; f. 8 *Cyp. lienardi*; f. 9 *Cyp. margarita*; f. 10 *Cyp. mariae*; f. 11 *Cyp. rabaulensis*; f. 12 *Cyp. katsuae*; figs. 13-15 *Cyp. saulae*; f. 16 *Cyp. hammondae*; f. 17 *Cyp. fimbriata*; figs. 18-19 *Cyp. minoridens*; f. 20 *Cyp. contaminata*; f. 21 *Cyp. ursellus*; f. 22 *Cyp. dillwyni*; f. 23 *Cyp. felina*; f. 24 *Cyp. pallidula*; f. 25 *Cyp. luchuana*; f. 26 *Cyp. martini*; f. 27 *Cyp. punctata*; f. 28 *Cyp. beckii*; figs. 29-30 *Cyp. kieneri*; f. 31 *Cyp. gracilis*; f. 32 *Cyp. hirundo*; f. 33 *Cyp. clandestina*; f. 34 *Cyp. asellus*.

Plate 10



Plate 11

