

SEA SHELLS
OF THE WORLD

BY

A. GORDON MELVIN, PH.D

Photographs by Lorna Strong Melvin

PAUL FLESCH & COMPANY

Melbourne & Sydney

Representatives

For Continental Europe:
BOXERBOOKS, INC., Zurich

For the British Isles:
PRENTICE-HALL INTERNATIONAL, INC., London

For Australasia:
PAUL FLESCH & CO., PTY. LTD., Melbourne

For Canada:
M. G. Hurtig Ltd., Edmonton

Published by the Charles E. Tuttle Company, Inc.
of Rutland, Vermont & Tokyo, Japan
with editorial offices at
Suido 1-chome, 2-6, Bunkyo-ku, Tokyo

Copyright in Japan, 1966
by Charles E. Tuttle Co., Inc.
All rights reserved

Library of Congress Catalog Card No. 65—18967

First Printing, 1966
Fourth printing, 1968

Book design by Simon Virgo
PRINTED IN JAPAN

Contents

INTRODUCTION · 7

SEA SHELLS OF THE WORLD · 13

GLOSSARY · 133

BOOK LIST · 137

INDEX · 141

Plate 21

1. *Conus sozoni* Bartsch. E.Mex. Yellow and brown markings and bandings. Sloped spire. Rare. See also Plate 17, No. 11.
2" 10.00
2. *Terebra dimidiata* L. Indo-Pac. Salmon mottling on white. Secondary band on each whorl. To 6".
3 1/4" 1.25
3. *Terebra duplicata* L. E.Afr. Alternately ribbed whorls of light grey and dark brown.
2 1/2" .60
4. *Terebra taurina* Sol. W.I. to Fla. Syn: *T. flammea* Lam. Cloudy orange markings on white. Florida specimens scarce and expensive.
3 3/4" 7.00
5. *Terebra chlorata* Lam. Indo-Pac. Two rows of blurred spots on each whorl.
2 3/4" .75
6. *Terebra muscaria* Lam. Indo-Pac. Three rows of square brown spots on each whorl, and four on body whorl. Secondary ring distinguishes this from *T. subulata* L.
4" 1.25
7. *Conus fulmen* Reeve. Japan. Brown mottling on faintly purplish white.
2 1/4" 1.50
8. *Conus zonatus* Hwass. Maldive Is. Dark bandings and markings, sometimes resembling tents. Rare. See also Plate 53, No. 19.
2" 26.00
9. *Conus vidua* Reeve. P.I. Widow Cone. White with dark mid and end bands. Tented. Rare form of *C. marmoreus* L.
2 1/2" 7.50
10. *Terebra strigata* Sow. Panamic Prov. Yellowish white, with strong, descending, dark stripes.
3 1/2" 1.50
11. *Terebra lanceata* L. Polynesia. White, with fine, descending, curved, brown lines.
1 3/4" 1.00
12. *Terebra pretiosa* Reeve. Japan. Long and slender. Sculptured in wide and narrow bands. Wider bands scalloped. 2 1/2" .75
13. *Conus granulatus* L. W.I. Reddish. Granulated surface with circular ridges. Indeterminate white bands. See also No. 18. Rare. 2 3/4" Imperfect specimens 5.00
14. *Terebra triseriata* Gray. Aust. Brownish. Many whorls. Delicately slender. Scarce.
3 1/2" 3.50
15. *Terebra strigilata* Chem. P.I. Shiny grey-brown. Whorls edged by black dots on white.
2 1/2" 1.00
16. *Terebra robusta* Hinds. Panamic Prov. Two or more rows of brown or black spots or stripes.
3" 1.25
17. *Conus vittatus* Hwass. Panamic Prov. Rib Bone Cone. Syn: *C. orion* Brod. Varies from white through orange and lilac to brown. Dark central band marked with white. Flame-shaped marks on spire. Not common.
1 1/2" 7.00
18. *Conus granulatus* L. W.I. Whitish specimen. See also No. 13.
19. *Terebra maculata* L. Indo. Pac. Marlin-Spike. Impressive and stocky. Brown squares and dots on cream. To 8".
5" 1.00
20. *Terebra guttata* Röd. P.I. Syn: *T. oculata* Lam. Salmon, with rings of white ovals. Incorrectly illustrated in some books. Rare.
3" 2.50
21. *Terebra diversa* Smith. Japan. Grey, with black squares on spiral white bands.
1 1/4" .50
22. *Terebra crenulata* L. Pac. Cream, ringed by brown dots and white knobs.
3" 1.00
23. *Terebra hoffmeyer* Abbott. P.I. A miniature. Shiny grey, with vertical ribs between whorls.
3/4" .50
24. *Terebra subulata* L. P.I. Cream, with two bars of squarish dark dots on edges of whorls. Distinguished by lack of secondary ring.
5" 1.50
25. *Terebra tigrina* Gmel. Indo-Pac. Syn: *Terebra felina* Dillwyn. Light color, with circling brown dots.
1 3/4" .75




Plate 48

1. *Murex elensis* Dall. W.Mex. Cream. Short spines. Lavender in aperture. $1\frac{1}{2}$ "—3". $1\frac{3}{4}$ " .60
2. *Leucozonia cingulata* Lam. Baja Calif. o Peru. Syn: *Opeatostoma pseudodon* Burrow. Dark rings on white. Perhaps longest apertural tooth of any gastropod. $1\frac{3}{4}$ " 1.00
3. *Cantharus mendozana* Berry, 1959. nΣ lege. W. Coast of Baja Calif. Named for collector Xavier Mendoza of Guaymas. White, with whorls ringed by pointed nodules. Scarce. $1\frac{3}{4}$ " 1.00
4. *Turris (Knefastia) tuberculifera* Brod. & Sow. W.Mex. Alternately banded with spiraling white nodules and dark brown bands. Rare. $1\frac{3}{4}$ "—2 $\frac{1}{4}$ ". 2" 2.00
5. *Distorsio decussatus* Val. W.Mex. Brown staining on white. Two folds on inner lip near top end of opening. $1\frac{3}{4}$ " 1.25
6. *Conus arcuatus* Brod. & Sow. W.Mex. Brown on white. High concave spire. $1\frac{1}{4}$ "— $1\frac{1}{2}$ ". $1\frac{1}{2}$ " 10.00
7. *Latirus mediamericus* Hertlein & Strong. W.Mex. Brown axial swellings. Not common. $1\frac{1}{2}$ " 1.75
8. *Phos articulatus* Hinds. W.Mex. Orange tints, with brown on axial knobs. $1\frac{1}{4}$ "— $1\frac{3}{4}$ ". $1\frac{1}{4}$ " 3.00
9. *Turris (Knefastia) walkeri* Berry. W. Mex. Brown. The nodal ribs cross spiral cords. $1\frac{1}{2}$ "— $1\frac{3}{4}$ ". $1\frac{3}{4}$ " 1.00
10. *Fusinus ambustus* Gould. W.Mex. Brown, with darker brown on axial knobs. $1\frac{1}{4}$ "— $1\frac{3}{4}$ ". $1\frac{1}{4}$ " .50
11. *Terebra formosa* Deshayes. W.Mex. White, with three bands of dots on body whorl. Rare. $1\frac{1}{2}$ "—3". $2\frac{1}{2}$ " 3.00
12. *Murex (Eupleura) muriciformis triquetra* Berry W.Mex. Greyish brown to white. Almost winged. Variable. $1\frac{1}{2}$ " 1.00
13. *Murex (Eupleura) muriciformis* Brod. W.Mex. $1\frac{1}{2}$ " .45
14. *Murex (Typhis) quadratus* Hinds. W. Mex. Light brown to white, with unique projecting tube on shoulder of body whorl. Rare. $\frac{3}{4}$ " 8.50
15. *Cancellaria urceolata* Hinds. W.Mex. Brown and cancellate. 1" 1.50
16. *Phos veraguensis* Hinds. W.Mex. Brownish and cross-ribbed. $1\frac{1}{8}$ " 2.00
- 17a. *Turris (Pleuroliria) oxytropis* Sow. W. Mex. Creamy white or buff. 1" 1.25
- 17b. *Turris (Pleuroliria) oxytropis albicarinata* Sow. W.Mex. Same as No. 17a, but buff with white keels. 1" 1.25
18. *Calliostoma bonita* Strong, Hanna, & Hertlein. W.Mex. Brown. Purple channel bordering inner lip. $\frac{1}{4}$ " 1.50
19. *Architectonica placentalis* Hinds. W. Mex. Brown and flat. Like Atlantic coast *A. peracuta* Dall. Rare. $\frac{1}{2}$ " 7.00
20. *Crucibulum serratum* Brod. W.Mex. White. Inner cup flat against shell. $\frac{1}{2}$ " 1.00
21. *Cosmioconcha palmeri* Dall. W.Mex. Shiny brown. Impressed line below suture. $\frac{3}{4}$ " 2.00
22. *Epitonium (Eglisia) nebulosa* Dall. W. Mex. Assigned tentatively to *Eglisia*. Brownish, with fine cancellation. No varices. $\frac{3}{4}$ " 6.00
23. *Natica broderipiana* Récluz. W.Mex. Brown, with three white bands which have brown spots. Not common. $1\frac{1}{2}$ " .80
24. *Fusinus panamensis* Dall. W.Mex. White. Pointed brown-tipped knobs. Rare. 4" 3.75
25. *Pleuroliria picta* Reeve, ex Beck Mss. W.Mex. White. Flecks of brown on spiral cords. 2" 1.25
26. *Sanguinolaria tellinoides* A. Adams. Panamic Prov. Brighter red within than without. $2\frac{1}{4}$ " 1.50
27. *Mitra hindsii* Reeve. W.Mex. Beautiful and delicate. Pink, with fine brown spiral lines. Not common. 1"— $1\frac{1}{2}$ " 3.00
28. *Polinices helicoides* Gray, 1825. Panamic Prov. Syn: *Natica glauca* Lesson, 1830. Thin and flat. Umbilicus overhung by a tongue. Not common. 2" 1.50

