



# PROCEEDINGS

OF THE

## Royal Society of Victoria.

VOL. XII. (NEW SERIES). 37111

PART II.

*Edited under the Authority of the Council.*

ISSUED APRIL, 1900.

*(Containing Papers read before the Society during the months of  
July, September, October, November, December 1899).*



THE AUTHORS OF THE SEVERAL PAPERS ARE SEVERALLY RESPONSIBLE FOR THE  
SOUNDNESS OF THE OPINIONS GIVEN AND FOR THE ACCURACY OF THE  
STATEMENTS MADE THEREIN.

MELBOURNE:

FORD & SON, PRINTERS, DRUMMOND STREET, CARLTON.

AGENTS TO THE SOCIETY:

WILLIAMS & NORGATE, 14 HENRIETTA STREET, COVENT GARDEN, LONDON.

To whom all communications for transmission to the Royal Society of Victoria,  
from all parts of Europe, should be sent.

1900.

AI

ART. XX.—*Catalogue of the Marine Shells of Victoria.*

PART III.

BY G. B. PRITCHARD AND J. H. GATLIFF.

[Read 12th October, 1899.]

In December, 1897, we contributed Part I. of the present catalogue, which included 83 species; and in December, 1898, Part II., which included 58 species; the present part includes 77 additional species, comprising the following families:—Pleurotomidæ, Conidæ, Strombidæ, Cypræidæ, Cassididæ, Naticidæ, Hipponycidæ, Capulidæ, Turritellidæ, and Vermetidæ. This brings the number of species we have already dealt with to 218.

Family PLEUROTOMIDÆ.

Genus *Drillia*, Gray, 1838.

*DRILLIA HARPULARIA*, Desmoulins.

*Pleurotoma harpula*, Valenciennes, m.s. (non Brocchi).

*Pleurotoma harpula*, Kiener. Icon. Coq. Viv., p. 58, pl. 18, f. 3.

*Drillia harpularia*, Desmoulins. Actes Soc. Linn. Bordeaux, vol. xii., p. 162.

1843. *Pleurotoma harpularia*, Reeve. Conch. Icon., vol. i., pl. 15, f. 124.

1884. *Drillia harpularia*, Tryon. Man. Conch., vol. vi., p. 193, pl. 14, f. 99.

1887. *Pleurotoma (Crassispira) harpularia*, Weinkauff. Conch. Cab. (ed. Küster) vol. iv., p. 97, No. 115, pl. 21, f. 2, a, b.

Hab.—Portland.

Obs.—This species has hitherto been somewhat rare amongst Victorian collections.

*DRILLIA QUOYI*, Desmoulins.

*Pleurotoma monile*, Valenciennes (non Brocchi).

Icon. Coq. Viv., Mon. *Pleurotoma*, p. 52, pl. 15, f. 3.

*Catalogue of the Marine Shells of Victoria.* 171

1842. *Pleurotoma quoyi*, Desmoulins. Actes Soc. Linn. Bordeaux, p. 61.

1846. *Pleurotoma quoyi*, Reeve. Conch. Icon., vol. i., pl. 16, f. 137.

1876. *Pleurotoma philipineri*, T. Woods. P.R.S. Tas., p. 136.

1884. *Surcula quoyi*, Tryon. Man. Conch., vol. vi., p. 242, pl. 7, f. 95.

1884. *Clavatula (Perrona) monile*, Tryon. *Id.*, p. 232, pl. 7, f. 96.

1884. *Pleurotoma philipineri*, Tryon. *Id.*, p. 167, pl. 34, f. 82.

1886. *Pleurotoma (Drillia) quoyi*, Watson. Chall. Zool., vol. xv., p. 304.

1887. *Pleurotoma (Surcula) quoyi*, Weinkauff. Conch. Cab. (ed. Küster) p. 101, sp. 121, pl. 22, f. 2.

1896. *Surcula quoyi*, Sowerby. P. Mal. S. Lond., p. 24.

Hab.—Western Port (dredged 5 fathoms, C. J. Gabriel); Apollo Bay, and Warrnambool (G. B. P.).

*DRILLIA BERAUDIANA*, Crosse.

1863. *Drillia beraudiana*, Crosse. Jour. d. Conch., p. 88, pl. 1, f. 6.

1884. *Drillia angasi*, Tryon (non Crosse). Man. Conch., vol. vi., p. 187, pl. 9, f. 36.

1887. *Pleurotoma (Crassispira) beraudiana*, Weinkauff. Conch. Cab. (ed. Küster), p. 95, No. 113, pl. 20, f. 7, 9.

Hab.—Flinders; San Remo; dredged off Phillip Island, Western Port, in about 5 fathoms (C. J. Gabriel); Port Phillip; Puebla coast.

*DRILLIA ÆMULA*, Angas.

1877. *Drillia æmula*, Angas. P.Z.S. Lond., p. 36, pl. 5, f. 9.

1880. *Drillia æmula*, Hutton. Man. N.Z. Moll., p. 44.

1884. *Drillia trailli*, Tryon (non Hutton). Man. Conch., vol. vi., p. 206, pl. 12, f. 37.

1887. *Pleurotoma (Drillia) æmula*, Weinkauff. Conch. Cab. (ed. Küster), p. 221, No. 259, pl. 41, f. 9.

1896. *Clathurella parvula*, Sowerby (non Reeve). P. Mal. S. Lond., vol. ii., p. 28.

Hab.—Port Phillip, Western Port; Portland (C. M. Maplestone); dredged off Phillip Island, Western Port, about 5 fathoms (C. J. Gabriel).

Obs.—The above species is fairly abundant in Port Phillip and Western Port. After carefully considering the meagre description and figure of *Pleurotoma parvula*, Reeve, Conch. Icon, vol. i., pl. 28, f. 254, and being unable to see the actual type we cannot agree with the conclusions of Sowerby in Proc. Mal. S. Lond., vol. ii., p. 28; our shell is more nearly related to *Clavatulæ spurca*, Hinds, and still closer to *C. rava*, Hinds.

The specimens dredged, above alluded to, are of greater length than usual, and the upper portion of the body whorl is not so angulate.

T. Woods himself acknowledges *Siphonalia pulchra* as an immature state of *Clathurella philomena*, in the Proceedings of the Royal Society of Tasmania for 1879, amongst some other corrections in a note to his paper on some Tasmanian Trochidæ.

*CLATHURELLA KYMATOËSSA*, Watson.

1886. *Drillia kymatoëssa*, Watson. Chall. Zool., vol. xv., p. 309, No. 39, pl. 26, f. 5.

Hab.—Off East Moncoeur Island, Bass Strait (Challenger).

Obs.—In general habit and sculpture this shell is very nearly related to *C. philomena*, T. Woods, and this is especially noticeable when it is compared with the large specimens of that species dredged in Western Port, probably a connecting series may be established.

*CLATHURELLA LEGRANDI*, Beddome.

1883. *Drillia legrandi*, Beddome. P.R.S. Tas., vol. 35, p. 167.

Hab.—Portland (C. M. Maplestone).

*CLATHURELLA ZONULATA*, Angas.

1867. *Clathurella zonulata*, Angas. P.Z.S. Lond., p. 113, pl. 13, f. 17.

1884. *Clathurella zonulata*, Tryon. Man. Conch., vol. vi. p. 285, pl. 17, f. 89.

Hab.—Western Port.

*CLATHURELLA BICOLOR*, Angas.

1871. *Clathurella bicolor*, Angas. P.Z.S. Lond., p. 18, pl. 1, f. 20.

1875. *Drillia atkinsoni*, T. Woods. P.R.S. Tas., p. 142.

1884. *Clathurella bicolor*, Tryon. Man. Conch., vol. vi., p. 284, pl. 16, f. 61.

Hab.—Western Port.

*CLATHURELLA SEXDENTATA*, Pritchard and Gatliff.

1899. *Clathurella sexdentata*, Pritchard and Gatliff. P.R.S. Vic., vol. xii, n.s., pt. 1. p. 104, pl. 8, f. 7.

Hab.—Sorrento Beach, Port Phillip.

Obs.—The type of this species is in Mr. Gatliff's private collection, and is a related form to *C. modesta*, Angas.

Genus *Raphitoma*, Bellardi, 1847.

*RAPHITOMA HARRISONI*, T. Woods.

1863. *Euryta pulchella*, Adams and Angas (non *Terebra pulchella*, Deshayes). P.Z.S. Lond., p. 418, pl. 37, f. 14.

1865. *Euryta pulchella*, Angas. *Id.*, p. 169.

1875. *Euryta brazieri*, Angas (non *Terebra brazieri*, Angas). *Id.*, p. 390, pl. 45, f. 5, 5a.

× 1877. *Mangelia harrisoni*, T. Woods. P.R.S. Vic., p. 56.

1884. *Daphnella harrisoni*, Tryon. Man. Conch., vol. vi., p. 306.

1885. *Terebra (Euryta) angasi*, Tryon. *Id.*, vol. vii., p. 38, pl. 12, f. 26.

1886. *Terebra (Euryta) angasi*, Tate. Southern Science Record, p. 4.

1887. *Pleurotoma (Cithara) harrisoni*, Gatliff. V.N., vol. iv., p. 59.

√ 1894. *Terebra (Euryta) harrisoni*, Brazier. P.L.S., N.S.W., vol. ix., p. 693.

Hab.—Western Port.