




Distribution of Shallow-Water
Marine Mollusca,
Yucatan Peninsula, Mexico

Harold E. Vokes
and
Emily H. Vokes

Mesoamerican Ecology Institute

Monograph 1



Middle American Research Institute

Tulane University, New Orleans

Publication 54. 1983

- Conus sennottorum* Rehder and Abbott, 1951 Pl. 20, fig. 8
Soc. Malac. "Carlos de la Torre," Revista, v. 8, p. 63, pl. 9, figs. 1, 2; Abbott, 1974, p. 255, fig. 2786
- Conus spurius atlanticus* Clench, 1942 Pl. 20, fig. 10
Johnsonia, v. 1, no. 6, p. 20, pl. 10, figs. 1-3; Abbott, 1974, p. 254, pl. 14, fig. 2779
- Conus villepini* Fischer and Bernardi, 1857 Pl. 20, fig. 11
Jour. de Conchyl., v. 5, p. 292, pl. 10 [not 9, as stated], fig. 12; Abbott, 1974, p. 256, pl. 14, fig. 2795

TEREBRIDAE

- Terebra (Terebra) dislocata* (Say, 1822) [*Cerithium*] Pl. 20, fig. 12
Acad. Nat. Sci. Phila., Jour., v. 2, pt. 2, p. 235; Abbott, 1974, p. 259, fig. 2836
- Terebra (Strioterebrum) protexta* (Conrad, 1846) [*Cerithium*] Pl. 20, fig. 13
Acad. Nat. Sci. Phila., Proc., v. 3, p. 26; Abbott, 1974, p. 260, fig. 2842
- Terebra (Hastula) cinerea* (Born, 1778) [*Buccinum*] Pl. 20, fig. 15
Index Mus. Caes. Vind., p. 262; 1780, Test. Mus. Caes. Vind., p. 267, pl. 10, figs. 11, 12; Abbott, 1974, p. 260, fig. 2847
- Terebra (Hastula) hastata* (Gmelin, 1791) [*Buccinum*] Pl. 20, fig. 14
Syst. Nat., ed. 13, v. 1, pt. 6, p. 3502; Abbott, 1974, p. 260, fig. 2846
- Terebra (Hastula) maryleeae* (Burch, 1965) [*Hastula*] Pl. 20, fig. 16
Veliger, v. 7, p. 242, pl. 31, fig. 4; Abbott, 1974, p. 260, fig. 2849
- Terebra (Hastula) sallaeana* Deshayes, 1859 Pl. 20, fig. 17
Zool. Soc. London, Proc., (1859), p. 287; Abbott, 1974, p. 260, fig. 2848

TURRIDAE

- Pyrgospira ostrearum* (Stearns, 1872) [*Drillia*] Pl. 21, fig. 1
Boston Soc. Nat. Hist., Proc., v. 15, p. 22; Abbott, 1974, p. 265, no. 2931
- Drillia coccinata* (Reeve, 1845) [*Pleurotoma*] Pl. 21, fig. 2
Conch. Icon., v. 1, *Pleurotoma*, pl. 33, fig. 299
- Neodrillia cydia* Bartsch, 1943 Pl. 21, fig. 3
Soc. Cubana Hist. Nat., Mem., v. 17, p. 84, pl. 7, fig. 6; pl. 10, fig. 3; pl. 14, fig. 1; Abbott, 1974, p. 71, fig. 3020
- Cerodrillia perryae* Bartsch and Rehder, 1939 Pl. 21, fig. 5
USNM Proc., v. 87, no. 3070, p. 128, pl. 17, fig. 1; Abbott, 1974, p. 271, fig. 3022
- Cerodrillia thea* (Dall, 1883) [*Drillia*] Pl. 21, fig. 4
USNM Proc., v. 6, [no. 384], p. 328, pl. 10, fig. 5; Abbott, 1974, p. 271, fig. 3023
- Crassispira (Crassispirella) cubana* (Melvill, 1923) [*Drillia*] Pl. 21, fig. 6
Malac. Soc. London, Proc., v. 15, p. 163, pl. 4, fig. 3; Abbott, 1974, p. 273, no. 3052, ? fig. 3071
- Crassispira (Crassispirella) fuscescens* (Reeve, 1843) [*Pleurotoma*] Pl. 21, fig. 7
Conch. Icon., v. 1, *Pleurotoma*, pl. 15, fig. 125
- Crassispira (Crassispirella) nigrescens* (C. B. Adams, 1845) [*Pleurotoma*] Pl. 21, fig. 8
Boston Soc. Nat. Hist., Proc., v. 2, p. 3; Abbott, 1974, p. 273, no. 3055
- Crassispira (Crassispirella) tampaensis* Bartsch and Rehder, 1939 Pl. 21, fig. 9
USNM Proc., v. 87, no. 3070, p. 136, pl. 17, figs. 5, 13; Abbott, 1974, p. 273, fig. 3054
- Pilsbryspira albomaculata* (Orbigny, 1842) [*Pleurotoma*] Pl. 21, fig. 10
in de la Sagre, Hist. l'Ile Cuba, Moll., Atlas, pl. 24, figs. 16-18; Abbott, 1974, p. 273, no. 3066
- Pilsbryspira harfordiana* (Reeve, 1843) [*Pleurotoma*] Pl. 21, fig. 11
Conch. Icon., v. 1, *Pleurotoma*, pl. 11, fig. 93
- Pilsbryspira jayana* (C. B. Adams, 1850) [*Pleurotoma*] Pl. 21, fig. 11
Cont. Conch., no. 4, p. 61
- Pilsbryspira leucocyma* (Dall, 1889) [*Drillia*] Pl. 21, fig. 11
USNM Proc., v. 6, [no. 384], p. 328, pl. 10, fig. 8; Abbott, 1974, p. 273, no. 3067
- Splendrillia (Syntomodrillia) janetae* (Bartsch, 1934) [*Leucosyrinx*] Pl. 21, fig. 11
Smithsonian Misc. Coll., v. 91, no. 2, p. 11, pl. 3, figs. 3, 11, 12; Abbott, 1974, p. 275, no. 3111
- Splendrillia (Syntomodrillia) moseri* (Dall, 1889) [*Drillia*] Pl. 21, fig. 11
Harvard Mus. Comp. Zool., Bull., v. 18, p. 97, pl. 36, fig. 3; Abbott, 1974, p. 275, fig. 3112

PLATE 20

CONIDAE, TEREBRIDAE

1. *Conus ermineus* Born
Height 32.5 mm
2. *Conus floridanus* Gabb
Height: a) 36.5 mm; b) 32 mm; c) 35 mm; d)
24.5 mm
3. *Conus granulatus* Linnaeus
Height 41.4 mm
4. *Conus jaspideus* Gmelin
Height 21 mm (W)
5. *Conus jaspideus stearnsii* Conrad
Height 24.5 mm (W)
6. *Conus jaspideus verrucosus* Hwass
Height 19.5 mm (W)
7. *Conus mus* Hwass
Height 24 mm
8. *Conus sennottorum* Rehder and Abbott
Height 35 mm (W)
9. *Conus regius* Gmelin
Height 46.5 mm
10. *Conus spurius atlanticus* Clench
Height 42 mm
11. *Conus villepinii* Fischer and Bernardi
Hypotype HMNS 6682, height 44.5 mm
12. *Terebra (Terebra) dislocata* (Say)
Height 51 mm
13. *Terebra (Strioterebrum) protexta* (Conrad)
Height 25 mm
14. *Terebra (Hastula) hastata* (Gmelin)
Height 23.2 mm
15. *Terebra (Hastula) cinerea* (Born)
Height 22.5 mm (W)
16. *Terebra (Hastula) maryleeae* (Burch)
Height 51 mm
17. *Terebra (Hastula) sallaeana* Deshayes
Height 51 mm




PLATE 20