

TRANSACTIONS OF THE  
WAGNER FREE INSTITUTE  
OF SCIENCE  
OF  
PHILADELPHIA

VOL. III.  
PART V.

DECEMBER, 1900

WAGNER FREE INSTITUTE OF SCIENCE  
MONTGOMERY AVE. AND SEVENTEENTH ST.  
PHILADELPHIA

CONTRIBUTIONS  
TO THE  
TERTIARY FAUNA OF FLORIDA

WITH ESPECIAL REFERENCE TO THE  
SILEX BEDS OF TAMPA AND THE PLIOCENE  
BEDS OF THE CALOOSAHATCHIE RIVER

INCLUDING IN MANY CASES  
A COMPLETE REVISION OF THE GENERIC GROUPS TREATED OF  
AND THEIR AMERICAN TERTIARY SPECIES

BY  
WILLIAM HEALEY DALL, A.M.  
PALEONTOLOGIST TO THE UNITED STATES GEOLOGICAL SURVEY; HONORARY PROFESSOR OF INVERTEBRATE  
PALEONTOLOGY IN WAGNER FREE INSTITUTE OF SCIENCE

PART V.  
TELEODESMACEA: SOLEN TO DIPLODONTA

PLATE XLI.

- Fig. 1. *Scala (Sthenorhytis) Mazyckii* Dall, basal view; Miocene of Cainhoy, South Carolina; diam. 28 mm. See *The Nautilus*, vol. ix., p. 111, February, 1896.
- Fig. 2. The same, front view.
- Fig. 3. *Ancillaria chipolana* Dall; Oligocene marl of the Chipola River, Florida; alt. 25 mm.
- Fig. 4. *Pleurotoma boadicea* Dall; Oligocene sands of Oak Grove, Florida; alt. 25 mm.
- Fig. 5. *Chama Willeoxi* Dall, interior of attached valve; Pliocene of Shell Creek, Florida; lon. 85 mm.
- Fig. 6. The same, with both valves in place.
- Fig. 7. The same, interior of upper valve; lon. 85 mm.
- Fig. 8. *Scala ranellina* Dall, basal view; Eocene of the Zeuglodon bed (Jacksonian), near Cocoa P. O., Alabama; diam. 23 mm. See *The Nautilus*, vol. ix., p. 111, February, 1896.
- Fig. 9. The same, front view; alt. (decollated) 33 mm.
- Fig. 10. *Astyris perforvida* Dall; Oligocene sands of Oak Grove, Florida; alt. 18 mm.
- ✕ Fig. 11. *Terebra psilis* Dall; Oligocene sands of Oak Grove, Florida; alt. 16.5 mm.

