

6. *Terebra albomarginata* Deshayes, 1859
 Solid, elongate with many whorls. Aperture almost rectangular, base of body whorl heeled. Pre-sutural groove deeply incised. Pre-sutural zone with a thick rounded raised band which is crossed by numerous low ribs, ribs continue as weak curved axial lirae on the anterior parts of the whorls, axial lirae crossed by strong spiral ribs forming a cancellate pattern. Whorls orange, pre-sutural zone white.

7 cm. Moderately uncommon. Indo-West Pacific; eastern Qld. as far south as Southport.

SEE ANANDA (3)
 GERNONHOESKY & BRATCHEV
 1976

ALBOMARGINATA (1)
 DESHAYES 1859
 = ANANDA (35)



212. *Terebra albomarginata*, Desh.

T. testa elongato-turrita, angusta, acuminata, aurantiaca, albo-marginata; anfractibus numerosis, angustis, planis, sulco impresso inaequaliter divisus, transversim striato-punctatis, striis quatuor; ultimo anfractu brevi, obtuso, canali brevi, contorto, angusto terminato; apertura minima, pallide lutea, ovato-subquadrata, angusta, extremitatibus attenuata; columella cylindracea, contorta, uniplicata.

Long. 44 mill., larg. 8.

Hab. L'Australie.

Collection Cuming.

Très-belle espèce rapprochée du *cingulifera* de Lamk., mais parfaitement distincte de ses congénères par sa coloration remarquable et les autres accidents de sa surface; le bourrelet blanc qui suit la suture est plissé et finement crénelé.

DESH 1859 p. 314

SEE ALSO: CERN 1966
 AT CINGULIFERA (2)

212. *T. albomarginata* Desh. P.Z.S., 1859, 314. Australia. Type B.M. A latticed shell. T 4a

11. *albomarginata*, *Terebra* - DESHAYES, 1859, P.Z.S.L., p. 314. Hab. L'Australie. Coll. Cuming. Size: 44 x 8 mm. Holotype: 44.5 mm (marked x inside aperture); this shell is yellowish-orange in colour with a white sutural band, obsolete axial plicae and 4 spiral rows of deep pits on each whorl. There is another specimen in the same box measuring 39.6 mm, and the loose label says "Coll. Dr. Gray" - no. 74.10.12.14. This Gray specimen is identical in sculpture with the holotype of *Terebra albomarginata*, except that it is uniformly dark brown in colour, and is the *T. cumingii* of authors, not of DESHAYES, 1857.

Terebra albomarginata Deshayes
 Thick, narrow shell with many whorls. Shoulder slopes gently just below suture and then whorls are almost straight-sided. Deeply incised pre-sutural groove with thick, rounded band above. Low ribs on pre-sutural band continued below as curved axial lirae, crossed by strong spiral ribs forming cancellation. Orange with pre-sutural zone white. Aperture orange. 7cm. Indo-Pacific (Pacific). Uncommon.

DANCE 77 p. 216

* (x) *T. (Dimidacus) albomarginata* DESHAYES, 1859: Proc. Zool. Soc. London, p. 314, no. 212; Fig.: REEVE; Conch. Icon., 1860, T., pl. 15, fig. 65. Apparently a rare species. Luzon. (DM no. 3539, in part). BURCH 1964



Species 65. (Mus. Cuming.)

Terebra albo-marginata. *Ter. testâ aculé turritâ, vividè aurantiâ, albo-marginatâ, anfractibus planis, superne sulco divisus, utrinque sulcum obliquè crenatis, deinde spiralliter liratis, interstitiis pertusis; apertura parvâ, columellâ contorto-recurvâ.*


THE WHITE-EDGED TEREBRA. Shell sharply turreted, bright-orange, edged with white, whorls flat, divided at the upper part by a groove, obliquely crenated on each side the groove, then spirally ridged, the interstices pricked; aperture small, columella twistedly recurved.

DESHAYES, Proc. Zool. Soc. 1859, p. 314.

Hab. Australia; Cuming.

Very conspicuous in respect of colour, the shell being bright-orange, except round the margin of the whorls where it is edged with white, and encircled throughout with ridges, the interstices between which are pricked.

RV 1860



Albomarginata (*Terebra*), Desh. Proc. Zool. Soc., 1859, p. 314, 29

T. ALBOMARGINATA, Desh. Pl. 9, fig. 54.

Whorls flat, bright orange, upper part white, divided by a groove, each side of which is crenated, below the groove spirally ridged, the interstices pricked. Length, 1.75 inches.

TRYON 1857 China.

54. *Terebra albomarginata*. Desh. Conch. Icon., f. 65.

1. *Terebra albomarginata*, Deshayes.

Terebra albomarginata, Deshayes, Review of *Terebra*, Proc. Zool. Soc. Lond., 1859, p. 314, sp. 212.

" " Reeve, Conch. Icon., vol. xii. pl. xv. fig. 65.
 " " Tryon, Manual, vol. vii. p. 29, pl. ix. fig. 24.

July 29, 1874. Levuka, Fiji. 12 fathoms.

Habitat.—Australia (Deshayes, British Museum, &c.)

WATSON 1886 p. 377

ALBOMARGINATA (2)

DESH 1859
= AMANDA (35)

ALLAN 196


White-edged Auger (*Perirhoe albomarginata*). Sharply turreted, whorls flat-tish, divided at upper part by a groove, obliquely crenulated on each side of groove, surface spirally ridged. Columella twisted and recurved. Bright orange, edged with white. Almost 2 inches long. Wide range throughout the Indo-Pacific as far north as China, and in recent years has been dredged in Sydney Harbour. Several rather similar species inhabit New South Wales and Queensland.

Many large Terebrids had been sorted out belonging superficially to *Perirhoe melamans* Iredale, but the keen eye of Mr. E. F. Nash divided them into three and his discrimination is correct.

The three forms referable to *Perirhoe* all have the lines punctate, and therefore belong to my subgenus *Dimidacus*; one of them agrees well with *Terebra albomarginata* Deshayes,⁶⁶ described from Australia, and not since recognized. Reeve's figure shows a young shell, but Hedley's MS. notes on the type in the British Museum (Natural History) agree with these specimens.

IREDALE 1931 p. 223

Fig. 4.—*Perirhoe albomarginata* Deshayes.



35a



4

35a, T.

Amanda Hinds. Holotype of *T. albomarginata* Deshayes, British Museum (N.H.) No. 197979; 44.5 mm. BRAT 87