

Pease's notes on Hawaiian species may be found in Amer. Journ. Conch. IV, pp. 123-125, and V, 86. There are no examples of his *Terebra assimilis* Pse. (= *T. contigua* Pse., 1871) in the Pease collection, Museum of Comparative Zoology. PILSBRY 1921-p300

T. ASSIMILIS, Pease.

T. aciculari, nitida; anfractibus laevis, longitudinaliter angulate plicato-costatis, superne, interstitiis punctatis; apertura oblongo-ovata; columella brevi, carinata, contorta; fusca, infra suturas grisea; columella aperturae fuscis.

Long. 15. Diam. 3 mill.

Hab. Insl. Oahu.

Shell acicular, shining; whorls flattened, smooth, longitudinally angulately plicately ribbed, interstices punctured above; aperture oblong ovate; columella short, keeled, twisted; brown, below the sutures greyish; columella and aperture brown.

PEASE 1869 p 67 V 15

26. ~~*assimilis*, *Acus* (*Abretia*) - ANGAS, 1867 P.Z. S. L. p. 11, pl. 13, fig. 8. Hab. Dredged in Port Jackson. Coll. Angas. Size: $5\frac{1}{2} \times 1\frac{1}{2}$ lines (= 12.4 x 3.4 mm). Holotype no. 70.10.26/53: 11.7 mm; syntype: 10.2 mm. = *Terebra fictilis* HINDS, 1844. CCG~~

PLACED HERE BY ERROR - SEE AT ASSIMILIS ANGAS 1867 PRECEDING PAGE.

Assimilis (*Terebra*), Pease. Am. Jour. Conch., v, p. 67, 1869. = *T. contigua*, Pease. Am. J. Conch., vii, p. 20, 39

TRYON 1880