

37. *booleyi*, *Terebra crenulata* var. — MELVILL & SYKES,  
 1898, P. M. S. L., 3: 44, pl. 3, fig. 5. Hab. Andaman  
 Islands. Size: not given. Holotype: 68.7 mm. =  
*Terebra crenulata* (LINNAEUS, 1758). C69


BOOLEYI  
 MELVILL & SYKES 1898

CRENULATA (11)

6. *T. crenulata* Lam. Melvill and Standen described a var. *booleyi*  
 in *Proc. Malac. Soc.*, iii, p. 42, pl. iii, f. 5, 1898, from the Andamans.  
 Type B.M. T44



11d,  
 Holotype of var. *booleyi* Melvill & Sykes, British Museum  
 (N.H.) No. 1898.4.40.1.; 68.7 mm. ERAT. 87



85. TEREBRA CRENULATA, L., var. BOOLEYI, n. var. (Pl. III, Fig. 5).  
 A very beautiful variety of this shell we had at first imagined to be  
 new. Warm fawn-colour, upon a flesh-coloured ground, its markings  
 are in longitudinal dashes and flames, irregularly disposed. But little  
 trace of crenulation is observable, the lower whorls being quite smooth,  
 though painted in a tessellate fashion between the sutural bands.  
 The upper whorls, however, are slightly noduled. n 85K 9814L ANDAMAN

5.  
*Terebra crenulata* var. *Booleyi*, n. var.