

154. *TEREBRA CÆLATA*, A. Adams et Reeve.

*Terebra cælata*, A. Adams et Reeve, Voy. du Samarang, p. 30. no. 3, pl. 10. f. 22.  
Hab. Philippines.

DESH 1859 p 304

CAELATA

A. ADAMS & REEVE  
1850

(52)

= FENESTRATA

44. *cælata*, *Terebra* - ADAMS & REEVE, 1850. Voy. Samarang, p. 30, pl. 10, fig. 22. Hab. China Seas. Size of figure: 40.6 mm. Holotype: 41.0 mm. = *Terebra fenestrata* HINDS, 1844.

C69

3. *TEREBRA CÆLATA*. Pl. X. Fig. 22. Ter. testâ lanceolato-turritâ, anfractibus longitudinaliter arcuatim plicato-costatis, costis liris duplicatis striisque transversis cancellatis, sulco unico conspicuo infra suturas; albâ.

HAB. Philippine Islands.

The chief peculiarity of this delicately carved species consists in the transverse ridges being finely duplicate.

ADAMS & REEVE 1850 p 30



22

12. *Terebra (Myurella) cælata*, Adams and Reeve. p 300

*Terebra cælata*, Adams and Reeve, "Samarang," p. 30, No. 3, pl. x. fig. 22.

" " Reeve, Conch. Icon., vol. xii. pl. xv. fig. 64.

" " Deshayes, Review of Terebra, Proc. Zool. Soc. Lond., 1859, p. 304.

" " Brazier, "Chevert" Exped. Shells, Proc. Linn. Soc. New South Wales, 1877, vol. i. p. 255, No. 32.

" " Tryon, Manual, vol. vii. p. 20, pl. v. fig. 77.

Station 203. October 31, 1874. Lat. 11° 6' N., long. 123° 9' E. Philippines. 20 fathoms. Mud.

*Habitat*.—China (British Museum); Philippines (Deshayes); Torres Strait, 20 fathoms (Brazier).

WATSON 1896 p 380



52e

52e, Middle whorls of questionable holotype of *T. cælata* Adams & Reeve.

32.—*TEREBRA (MYURELLA) CÆLATA*.

*Terebra cælata*, Adams and Reeve, Moll. Voyage, Samarang, p. 30, pl. 10, f. 22.

" " Reeve, Conch. Icon., pl. 15, sp. 64.

Hab. Darnley Island, Torres Straits, 20 fathoms, sandy mud bottom.

BRAZIER 77 1877

*Cælata* (*Terebra*), Adams and Reeve. Moll. Voy. Samar., 30, pl. x. f. 22, . . . . . 20

*T. CÆLATA*, Adams and Reeve. Pl. 5, fig. 77.


Surface sharply granulated by longitudinal and spiral sculpture; transparent white. Length, 41 mill.

Has the form of the preceding species, from which it is, perhaps, insufficiently distinguished by the non-development of the inferior band of tubercles.

TEXTILIS

China.

TRYON 1878 p 20



77 *T. cælata*  
Adams & Reeve.

17. *Terebra cælata*, Ads. and Reeve. Reeve, Icon., f. 64

Species 64. (Mus. Cuming.)

*TEREBRA CÆLATA*. Ter. testâ acutè tereti, pellucido-albâ, anfractibus plano-convexis, supernè sulco divisis, utrinque sulcum obliquè granatis, undique profundè clathrato-cælatis; aperturâ parvâ, columellâ contorto-recurvâ.


THE CARVED *TEREBRA*. Shell sharply terete, transparent-white, whorls flatly convex, divided by a groove at the upper part, obliquely grained on each side the groove, deeply lattice-carved throughout; aperture small, columella twistedly recurved.

ADAMS and REEVE, Moll. Voy. Samarang, p. 30. pl. 10. f. 22.

Hab. China; Belcher.

Remarkable for the high relief in which the elaborate lattice-work of this species is carved throughout.

RV 1860



64.