

91. *Terebra caliginosa* Deshayes, 1859

(Pl. 24, figs. 91a-b)

1859 *Terebra caliginosa* Deshayes, Proc. Zool. Soc. Lond. pt. 27: 287; 1860 Reeve, Conch. Icon. 12, pl. 20, fig. 100.

1978 *Hastula caliginosa* (Deshayes), Cernohorsky, Trop. Pacific Mar. Shells, p. 150, pl. 53, fig. 8.

Description: Shell to 30 mm; color dull purplish brown or purplish gray; outline of whorls almost straight except for convex body whorl; subsutural band absent except for grooves between ribs; axial ribs narrow, widely-spaced, straight, becoming sinuous on body whorl; spiral sculpture absent; sutures shallow; body whorl extremely elongate, with ribs fading at the periphery; aperture semi-elongate, with brown interior; columella stained brown, curved, with slight parietal callus; siphonal fasciole with cord-like keel.


Type locality: "Philippine Islands."

Distribution: From Durban, South Africa to the Philippines.


Type: Holotype BM(NH) no. 197987; 29.7 × 6 mm.

Discussion: This is the type species of the genus *Punctoterebra* Bartsch, 1923. It may be separated from *T. nitida* (92) by its brown columella and interior of aperture, lack of shine, and very shallow, almost unnoticeable subsutural grooves.

8-91-197



91a,b, *T. caliginosa*
Deshayes. 91a, Holotype British Museum (N.H.) No. 197987;
29.7 mm. 91b, Fiji Is.; 28.0 mm.



62. *TEREBRA CALIGINOSA*, Desh.
T. testa elongato-subulata, angusta, castaneo-grisea, livida, longitudinaliter regulariterque acute costata; anfractibus planis, conjunctis, linea impressa vix perspicua, inaequaliter divisis, interstitiis costularum obsoletissime transversim striatis; ultimo anfractu elongato; apertura minima, ovato-attenuata, fusca, canali brevi latoque terminata; columella cylindracea, extus angulo acuto marginata.
 Long. 30 mill., larg. 6.
 Hab. Isles Philippines.
 Collection Cuming.


DESH 1859 p 287

46. *caliginosa*, *Terebra* - DESHAYES, 1859, P. Z. S. L., p. 287. Hab. Isles Philippines. Coll. Cuming. Size: 30 x 6 mm. Holotype: 29.7 mm. C69

62. *T. caliginosa* Desh. P.Z.S., 1859, 287. Philippines. Type B.M. J. 44

*(x) *T. (? Hastula) caliginosa* DESHAYES, 1859: Proc. Zool Soc. London, p. 287, no. 62; Fig.: REEVE; Conch. Icon., 1860, T., pl. 20, no. 100. Has not been among specimens examined. BUZE464

Olive gray; axial ribs crisp and narrow; to approx. 30 mm. (BMNH type collection)



Hastula caliginosa (Deshayes)
 NEW BRITAIN-PHILIPPINES SK

8. *Hastula caliginosa* (Deshayes). West Pacific. To 1 1/4 inches. P. 150

Hastula caliginosa (Deshayes, 1859)

Plate 53 Figure 8

Shell about 25-30mm in length, shining, whorls flat, sutures wavy, sculptured with strong, evenly angulate axial ribs, ribs becoming obsolete on the dorsal side of the body whorl; presutural area with a row of deep pits between the axial ribs, interspaces of ribs with obsolete, fine spiral striae which are discernible only under magnification, columella with a basal fold, siphonal fasciole with a prominent, oblique cord. Uniformly greyish-brown or lead-grey in colour, early whorls usually paler.

Moderately rare. From New Britain westward to the Philippine Islands; subtidal. C 784150

Species 100. (Mus. Cuming.)

TEREBRA CALIGINOSA. *Ter. testā gracili-subulatā, saturatē livido-purpureā, anfractibus declivi-planatis, indivisis, longitudinaliter varicoso-costatis; apertura parvā, columellā ad basin subrecurvā.*

THE DARK TEREBRA. Shell slenderly subulate, dark livid-purple, whorls slopingly flattened, undivided, longitudinally varicosely ribbed; aperture small, columella slightly recurved at the base.

DESHAYES, Proc. Zool. Soc. 1859, p. 287.

Hab. Philippine Islands; Cuming.

This species is founded on a single specimen in Mr. Cuming's collection, of a dark livid-olive or lead-colour.



RV 1800



11. *Hastula caliginosa* (Deshayes, 1859)
 38.2mm; Cebu.
 Shell small to medium sized, glossy, with a tall spire and moderately impressed sutures; sculptured with rounded axial ribs, smooth between, and a punctate presutural cord; aperture elliptically ovate, columella folded; totally grey in colour, aperture orange-brown; reaches 40mm.
 DISTRIBUTION: Sporadically found throughout the Philippines. SPRIU & LLOYD 1966 p 260

Caliginosa (*Terebra*), Desh. Proc. Zool. Soc., 1859, p. 287, 35

T. CALIGINOSA, Desh. Pl. 10, fig. 97.
 Shell slender, varicosely ribbed; dark chocolate.
 Length, 1.25 inches.

Philippines.

Founded on a single specimen in the Cumingian Collection. It may = the preceding, but punctured interspaces are neither figured nor described.

Dr. von Martens refers here the unfigured *T. Kirki*, Hutton, of New Zealand. TRYON 85 p 31



97
T. caliginosa
 Desh.

97. *Terebra caliginosa*, Desh. Reeve, Conch. Icon., f. 100