

Whitish with pale brown markings; to 29 mm. (BMNH type coll.)

Terebra cinctella Deshayes
PAKISTAN S.K.

158. TEREBRA CINCTELLA, Desh.

T. testa elongato-angusta, subulata, longitudinaliter costellata, griseo-fusca, livida, nitida; anfractibus convexiusculis, sulco lato inaequaliter bipartitis, transversim striatis, striis quatuor vel quinque impressis; ultimo anfractu brevi, fasciola pallida cincto; apertura minima, ovato-angusta, extremitatibus attenuata, intus fusca; columella brevi, cylindracea, basi canali brevi terminata.

Long. 29 mill., larg. 6.
Hab. L'embouchure de l'Indus.
Collection Cuming.

1859 | 305 DESH

51. *cinctella*, *Terebra* - DESHAYES, 1859, P. Z. S. L., p. 305. Hab. L'embouchure de l'Indus. Coll. Cuming. Size: 29 x 6 mm. Holotype: 27.8 mm; syntypes: 31.0 mm (erroneously marked x inside aperture), and 26.9 mm.

158. *T. cinctella* Desh. P.Z.S., 1859, 305. Mouth of the Indus. Types (3) B.M. T44

TEREBRA CINCTELLA Desh. (= *undulata* Gray).

I. Bombay (*Abercrombie*). Karachi (*Mus. Brit.*). 1745 1901 1428

158. *T. cinctella*, Deshayes, variety of *T. picta*, Hinds. 2011600

Cinctella (*Terebra*), Desh. Zool. Proc., 305, 1859.

= *T. undulata*, Gray, 23

T440N1885

96. *Terebra cinctella* Deshayes, 1859

(Pl. 26, figs. 96a-b)
(Color pl. D, fig. 23)

1859 *Terebra cinctella* Deshayes, Proc. Zool. Soc. Lond. pt. 27: 305.

Description: Shell to 31 mm; color fawn to amber, some with lighter subsutural band and light peripheral stripe on body whorl which shows through outer lip, others with band of lavender hue; outline of whorls convex; protoconch of 2 1/2 small whorls; subsutural band convex, moderately narrow, with elongate nodes, defined by deep suture and constricting subsutural groove, usually with punctations between ribs; axial ribs curved, evenly spaced, well-developed, contiguous with elongate nodes on band; interspaces about equal to ribs; spiral cords numerous, fine, 6 to 12 on penultimate whorl, sometimes also appearing on band; aperture moderately quadrate; columella recurved.


Type locality: "Mouth of the Indus," Pakistan.

Distribution: From the Arabian Sea to the Solomon Is.

Type: Lectotype BM(NH) no. 197988; 27.8 x 5.8 mm.

Discussion: A good diagnostic feature of this species is the constriction both above and below the narrow, convex subsutural band. *T. swainsoni* Deshayes lacks this.

B-96-100



FIGS. 96a, b, *Terebra cinctella* Deshayes. 96a Lectotype British Museum (N.H.) No. 197988; 27.8 mm. 96b, Paralectotype, same Institution No. 197988; 31.0 mm.