

261. *Duplicaria albozonata* (E. A.

Smith, 1875)

(Pl. 67, fig. 261)

1875 *Terebra albozonata* E. A. Smith, Ann. & Mag. Nat. Hist. ser. 4, 15:415; 1879 E. A. Smith, Proc. Zool. Soc. Lond. p. 185, pl. 19, fig. 5.


**Description:** Shell to 25 mm; color pale brown with a narrow white zone above the suture and at the periphery of the body whorl; outline of whorls almost flat except for inflated body whorl; subsutural band defined by narrow deep groove; axial ribs, narrower than interspaces, spiral sculpture absent; aperture broad, quadrate; columella recurved.

**Type locality:** "Matoza Harbour, Japan, 6 fms."

**Distribution:** Japan.

**Type:** Lectotype BM(NH) no. 1873.8.6.9; 24.7 × 8.0 mm.

**Discussion:** The two specimens in the type lot are worn and faded. The lectotype has a bad break on the body whorl which causes it to look more inflated than it might be if it were not damaged. The paralectotype is a very young specimen of 8 mm. We have not seen any recently collected specimens which would match the type of *T. albozonata*. B.261 - b 212



261. *D. albozonata* (E. A. Smith). Lectotype British Museum (N.H.) No. 1873.8.6.9.; 24.7 mm.



8. TEREBRA ALBOZONATA, Smith. (Plate XIX. fig. 5.)  
*Terebra albozonata*, Smith, Ann. & Mag. Nat. Hist. 1875, vol. xv. p. 415; l. c. 1877, vol. xix. p. 226.  
*Hab.* Station 27. Matoza Harbour (l. c.).  
It is satisfactory to have obtained a second, although young, example of this species agreeing perfectly with the type.

SMITH 1875 P256 p 185

*Terebra albozonata*, sp. nov.

Ann. & Mag. Nat. Hist. 1875, vol. xv. p. 415.

Testa subulata, dilute fusca, zona alba angusta ad anfractuum basim cincta, et circa peripheriam anfr. ultimi zona alba ornata; anfract. 12, primi 2 laeves, convexi, caeteri fere plani, costis longitudinalibus numerosis, gracilibus, obliquis (in anfr. ult. 19-20 versus basim attenuantibus) instructi, incrementi lineis striati, et sulco unduloso spirali inaequaliter divisi, anfr. ultimus rotundatus; columella brevis; canalis brevis latus paululum recurvus.  
Long. 25 mill., diam. 7.

*Hab.* Matoza Harbour, Japan (Commander St. John).

This species probably attains a much larger size than the above dimensions. The palish brown colour, with the narrow white zone at the base of the whorls and that around the periphery, and the numerous oblique ribs, which are divided above by a fine furrow, are the chief peculiarities. The nearest ally appears to be *T. badia*, Desh.

E.A. SMITH 1877 p 226

- 12. *albozonata*, *Terebra* - E. A. SMITH, 1875, A. M. N. H., 15: 415 and 1877, A. M. N. H., 19: 226. *Hab.* Matoza Harbour, Japan (Commander St. John). Coll. J. G. Jeffreys. Size: 25x7 mm. Holotype, no. 73.8.6.9: 24.7 mm (spot of red wax on labial lip); paratype: 8.0 mm.

C64

*Terebra albozonata* Smith.

Proc. Zool. Soc. 1879. pag. 185. Tab. 19. fig. 5.

Species fusca taenia alba cincta.

Kiushiu.

DONKER 1881 p 73

~~SMITH 1877 p 226~~



- 6. *Diplomeriza albozonata* (Smith) • Height 57 mm.; diam. maj. 13 mm., min. 11.8 mm.

BRADSHAW 34



- 16. *Diplomeriza albozonata* (Smith). シロオビトクサ H. 57, Dj. 13, Dn. 11.8. Loc. Wakayama-ken.

SMITH 1875 p 415 JAPAN

4. *Terebra albozonata*\*, sp. nov.  
*Hab.* Matoza Harbour, 6 fathoms, sand.  
This species is of a pale brown colour, with a narrow white zone above the suture, and also one around the periphery; it is furnished with numerous oblique ribs, which are cut across by a fine infrasutural furrow.