

Terebra coerulescens var. T. castanea Kiener p. 19. pl. 7. Fig. 14 Ile de France.

NOTES 1880 p. 230

CASTANEA (Kiener)

HINDS 1844

= BADIA (248)

Castanea (Terebra), Hinds (not Kiener). Thes. Con., i, p. 161, pl. 43, f. 59. = T. badia, Desh. 15

T. Kiener 1880

32. TEREBRA CASTANEA, Kiener, l. c. p. 19. t. 7. f. 14.

Hab. Mauritius: Kiener. Guinea: Humphrey. (v. s. in m. Cuming.)

I feel tolerably certain that I have referred the right shell to Kiener's description and figure. With the latter it agrees sufficiently well, but the whorls are decidedly divided. His description and figures widely differ.

H 444 p 162

31. T. CASTANEA. (pl. xliii. f. 59.) Kiener, l. c. p. 19. t. 7. f. 14.

Testâ elongatè conico-subulatâ, subturritâ, nitidâ, castaneâ; anfractibus planulatis, sulco impresso divisis, longitrorsùm acutèplicatis; interstitiis lævigatis; aperturâ supernè attenuatâ; columellâ nudâ, rectiusculâ, subtruncatâ; canali nullo.

The folds are separated by distinct smooth spaces, and are sharp and nearly straight; the girdling is effected more by the cessation of the folds, than by any constriction. The colour is a very dark chesnut, with sometimes a slight appearance of white near the sutures; the last whorl is banded by white, and its base is somewhat tubercular.

Mauritius: Kiener. Guinea: Humphreys.

H. 445. T. E. p. 161 t. xliii f. 59.

