

238. *Duplicaria concolor* (E. A. Smith,  
1873)

(Pl. 61, fig. 238)

1873 *Terebra concolor* E. A. Smith, Ann. & Mag. Nat. Hist., ser. 4, 11:65; 1966 Cernohorsky & Jennings, Veliger 9(1):62, pl. 6, fig. 45 (illustrated holotype).


**Description:** Shell to 22 mm; color whitish; outline of whorls slightly convex; subsutural band moderately broad, axially ribbed, defined by narrow groove; axial ribs narrower than interspaces, about 18 on penultimate whorl; spiral sculpture absent; aperture quadrate; columella almost straight.

**Type locality:** None given.

**Distribution:** Unknown.

**Type:** Holotype BM(NH) no. 1858.5.12.252; 22.0 × 6 mm.

**Discussion:** No specimens have been found to match the holotype, which is in poor condition. This may prove to be a juvenile specimen of *D. duplicata* (237).  
E. 238-198



238. *D.*

*concolor* (E. A. Smith). Holotype British Museum (N.H.)  
No. 1858.5.12.252.; 22.0 mm.

*Duplicaria concolor* (E. A. Smith, 1873)

(Plate 53. Figure 394.)

**SHELL:** Shining-white in colour, sculptured with a prominent spiral ledge at sutures and about 14 wide-spaced axial ribs; the presutural ledge separates long and oblique nodules from the rest of the whorl.

**SIZE:**  $\frac{1}{4}$  to one inch.

**DISTRIBUTION:** West Pacific. Rare.

**REMARKS:** Illustrated is the lectotype of *D. concolor* preserved in the British Museum (Nat. Hist.) No. B.M.N.H. 1965136 (Length: 22.0 mm).

CONCOLOR (2)

E.A. SMITH 1873

(V)

(238)



C\_1967

Figure 45: *Duplicaria concolor* (E. A. SMITH). Lectotype no. B. M.N.H. 1965136. Length 22.0 mm. Photo British Museum (Natural History)

C&J 1966 Top F.

See also C&J Top F.

DUPICARIA SPEC - p 98.  
fig 44.

56. *concolor*, *Terebra* - E. A. SMITH, 1873, A. M. N. H., 11: 265. Hab. ? Size: 22x6 mm. Holotype no. 1858. 5.12.252: 22.0mm. This appears to be a worn specimen of *Duplicaria jukesii* (DESHAYES, 1857). C69



45



x1



x4

Shell white with low, rather widely separated axial ribs; to approx. 25 mm. (BMNH type collection)

*Terebra concolor* E.A. Smith

WESTERN PACIFIC

S.K.

*Concolor* (*Terebra*), E. A. Smith. Ann. Mag. N. H., 1873, xi, p. 265, . . . . .

39

TR40N 1825

*Terebra concolor*.

*T.* testa polita, nitente, aut alba aut dilutissime caeruleo-alba, subulata, subturrita; anfr. planiusculi, sulco bene definito inaequaliter divisi; pars superior nodulis parvis, elongatis, obliquis munita; pars inferior costis parvulis, subdistantibus, erectis (in anfr. ultimo breviusculo 14-15 ad basim sensim obsolete) instructa; columella curta, subrecta; canalis brevissimus.

Long. 22 mill., diam. 6; apertura long. 5 mill., diam. 2 $\frac{1}{2}$ .

*Hab.* — ?

A whitish shining species, furnished with oblique oblong nodules on the infrasutural belt; and on the remainder of the whorls there are small, regular, upright ribs somewhat distant from each other, which gradually become stouter towards the apex.

SMITH 1873 p 265