

T. COSTELLIFERA, Pease, 1869

T. subulata, spira gracilis; anfr. plano-convexis, laevibus, longitudinaliter forte angulate plicato-costatis, infra suturam profunde sulcatis; sutura, sulcoque subangulatis; apertura ovata; columella contorto-recurva, laminata; albida, anfr. partem inferiorem cinereis, infra suturam purpurascete cinerea maculosis; apice interdum fulvescente.

Long. 20. Diam. 5 mill.

Hab. Insl. Oahu.

Shell subulate, spire slender; whorls flatly convex, smooth, longitudinally strongly angularly plicately ribbed; below the suture deeply grooved; suture and edge of groove slightly angulate; aperture ovate; columella twistedly recurved, laminate. Whitish, lower part of the whorls pale ash color, blotched or spotted below the suture with purplish ash color, apex somewhat yellowish.

The specimens of this species collected do not appear to be mature.

PEASE 1869 p 66

COSTELLIFERA  
PEASE 1869

=THAANUMI 252

Additional species have been recorded,


but the records are either unconfirmed or erroneous:

T. costellifera Pease, 1869b (not identifiable from the original description);

AK 79 p 403

Costellifera (Terebra), Pease. Am. Jour. Conch., v, p. 66, 1869, (39)

TRAYON 1885



252b

BAIT 87 252b. Immature lectotype of T. costellifera Pease. Museum Comparative Zoology, Harvard No. 148800; 19.3 mm.