

199. *Terebra fatua* Hinds, 1844

(Pl. 51, figs. 199a-c)

(Color pl. E, fig. 14)

1844 *Terebra fatua* Hinds, Proc. Zool. Soc. Lond. pt. 11:150; 1844 Hinds in Sowerby, Thes. Conch. 1:158, pl. 42, fig. 28; 1983 Bouchet, Boll. Malac. 18(9-12):194, figs. 36-37.

1857 *Terebra pura* Deshayes, Proc. Zool. Soc. Lond. pt. 27:82, pl. 5, fig. 8.

**Description:** Shell to 75 mm; color, all creamy white or with pale orange-brown spots anterior to suture and at periphery of body whorl; outline of whorls straight; subsutural band confined to early whorls on some individuals, weakly indicated throughout or entirely absent in others; axial sculpture of straight ribs resembling those of a *Duplicaria* on early whorls, often fading completely after about 7 whorls of the teleoconch with only axial striae remaining; body whorl exceptionally long; aperture moderately elongate; columella straight.


**Type locality:** *T. fatua*, "St. Christopher, W. Indies," error. Boavista, Cape Verde Islands, is here designated as the type locality; *T. pura*, "Zanzibar," error.

**Distribution:** Endemic to the Cape Verde Islands; 2-15 m.

**Type:** *T. fatua* lectotype BM(NH) no. 1968175, 74.9 mm; *T. pura* lectotype BM(NH) no. 197970, 64.7 mm.

**Discussion:** This rare species has an extremely long body whorl. It has not been reported from the Western Atlantic subsequent to its original description, and it may be assumed that the species is endemic to the Cape Verde Islands.

B-199-1169



14: *Terebra fatua* Hinds; Cape Verde Is.; 38.4 mm.

199a-c, *T. fatua* Hinds. 199a, Lectotype British Museum (N.H.) No. 1968175; 74.9 mm.

199c, Boavista Id., Cape Verde Ids.; 38.4 mm.

*Terebra fatua* HINDS, 1844

Figs. 4, 36-37

BOUCHER 1983 p

*Terebra fatua* HINDS, 1844a: 150; 1844b: 158, pl. 42 fig. 28.


**Localité type:** la localité-type indiquée par HINDS (« St. Christopher, West Indies ») est erronée.

**Matériel type:** 2 syntypes BMNH 1968175.

**Matériel examiné:** le matériel type et environ 25 individus et coqs. des îles du Cap-Vert, essentiellement Boavista. 2-4 m (coll. Rolan, Ramalho, Soares, Saunders) et CALYPSO st 76, 78, devant Praia Colhino 10-15 m.

**Distribution:** endémique aux îles du Cap-Vert.

**Remarques:** *Terebra fatua* est une espèce à larves non planctoniques descendant du même ancêtre pliocène que *T. senegalensis* (BOUCHER 1981 sous le nom de *Terebra* n. sp.). Il se distingue de *T. senegalensis* et *T. concinna* par sa couleur rose crème unie, sans taches, y compris dans la bande sous-suturale. La sculpture axiale est présente sur les tours jeunes, puis disparaît sur les tours adultes.



FATUA (2)

HINDS 1844

(199)  
V

36 et 37, *T. fatua*, Sal Rei, Boavista, Îles du Cap Vert, 2-4 m; 33 mm et 63 mm.

74. *fatua*, *Terebra* - HINDS, 1844, P.Z. S. L., p. 150. Hab. St. Christopher, West Indies (Mr. Miller, 1799). Coll. Cuming. Size: 34 lin. (= 76.5 mm). Holotype no. 1968175: 74.9 mm (marked x inside aperture); syntype: 62.7 mm. C69

TEREBRA FATUA. *Ter. testâ turrito-subulatâ, albidâ, lævigatâ, politâ; anfractibus subplanulatis, superioribus lineâ impressâ cinctis, maculis fuscis pallidis distantibus biseriatis ornatis; spirâ obsolete plicatâ; anfractu ultimo elongato, maculis exceptis, unicolore.* Axis 34 lin.

Hab. St. Christopher, West Indies; on the sand: Mr. Miller, 1799.

Cab. Cuming.

H. 1844 b T. c. p. 158 pl. 42 f. 28

20. *T. FATUA*. (pl. xlii. f. 28.) *Hinds*, l. c. p. 150.

Testâ turrito-subulatâ, albidâ, lævigatâ, politâ; anfractibus subplanulatis, superioribus lineâ impressâ cinctis, maculis fuscis pallidis distantibus biseriatis ornatis; spirâ obsolete plicatâ; anfractu ultimo elongato, maculis exceptis, unicolore.

Shell subulate, subturritid, white, smooth; whorls flat, lengthened, polished, the upper girdled, ornamented on the margins with two series of very pale somewhat distant brown spots; last whorl elongated, those of the spire plicate. Aperture elongated, effuse. Columella straight, truncate; no canal.

St. Christopher, West Indies, on the sand: Miller, 1799.

H. 1844 b T. c. p. 158 pl. 42 f. 28



16. *TEREBRA FATUA*, Hinds.

*Terebra fatua*, Hinds, *Thes. Conch.* p. 58. no. 20. pl. 42. f. 28.

Hab. St. Christophe (Indes occidentales). DESM 1859 h 298

16. *T. fatua*, Hinds, a bleached state of *T. cingula*, Kiener. RV 1860b

*Fatua* (*Terebra*), Hinds. *Proc. Zool. Soc.*, 1843, p. 150.  
= *T. cingula*, Kien. 1843



TRAYON 1875

28. *Terebra fatua*, Hinds (= *cingula*). Sowb., *Thes. Conch.*, f. 28,