

E[pitonium]feldmanni RÖDING in Bolten, Mus. Boltenianum, p. 94, 1798. Ref. to "Gmel. Buccin. subulatum sp. 131. 7. Martini 4. t. 154 1446. 2 St." [No locality cited by Chemnitz.] HEDLEY, Proc. Linn. Soc. New South Wales, vol. 38, pt. 2, p. 306, 1913. "It does not appear to have been remarked that *Epitonium feldmanni* Bolten, 1798, is an earlier name for Lamarck's species" [that is, *Terebra flammea* Lamarck].

Clench considered *Terebra feldmanni* to be close to or identical with *T. robusta*. Upon comparison of figure 1446, given by Chemnitz in the Conchylien-Cabinet, with the original figure of *T. robusta* we fail to see sufficient similarity to warrant relegating this well known west American name to the synonymy of *T. feldmanni*. We are inclined to agree with Reeve, Hedley, and Abbott in considering *T. feldmanni* to be a Caribbean species.

H & W 1361 b 79

FELDMANNI

RÖDING 1798

= TAURINA

(42)

1216/40 *E. feldmanni* Die Bildmanche Penne,
Gmel, Buccin. subulatum sp. 131. 7.
Martini 4. t. 154 1446. 2 St

RÖD 1798 194