

FLAVA
GRAY 1834
65
= AFFINIS

TEREBRA FLAVA. *Ter. testâ turrilâ, ovato-subulatâ, longitudinaliter plicatâ plicis frequentibus aequalibus, striis spiralibus frequentibus punctatis; anfractibus planis, cingulo convexiusculo; pallidè flavâ.*
G. 34; 60

Axis 1 unc.
Far. Subulata, gracilis, costis longitudinalibus magis acutis.

Species 126. (Mus. Gray.)

TEREBRA FLAVA. *Ter. testâ conico-subulatâ, flavescente-albâ, anfractibus plano-convexis, sulco pertuso supernè divisis, longitudinaliter plicato-liratis, interstitiis lineis spiralibus impressis decussatis; aperturâ parvâ, columellâ ad basin productâ et paululum recurvâ.*

THE YELLOW TEREBRA. Shell conically subulate, yellowish-white, whorls flatly convex, divided by a punctured groove at the upper part, longitudinally plicately ridged, interstices decussated by impressed spiral lines; aperture small, columella produced at the base, and a little recurved.

GRAY, Proc. Zool. Soc. 1834, p. 60.
Hab. —?



A yellowish conoid shell, longitudinally ridged, with the interstices closely linearly grooved. It is scarcely mature, and not in very good condition. The species must be regarded as doubtful. RJ 1860

77. T. FLAVA. (pl. xlv. f. 75.) Gray, l. c. p. 60.

Testâ obeso-subulatâ, supernè attenuatâ, subturritâ, pallidè flavâ, anfractibus planulatis, supernè lineâ impressâ divisis, longitrorsum plicatis, striis transversis cancellatis; plicis subrectis, confertis; aperturâ abbreviatâ; columellâ nudâ, contortâ.

Shell not uniformly subulate, the superior whorls being proportionately more so than the others; this with the flattened cancellated whorls and uniform pale yellow colour will readily distinguish it. Native country unknown.

H 446 T.C. p 177 6 xlv f 75.



81. TEREBRA FLAVA, Gray, l. c. p. 60.
Hab. —? (v. s. in m. Gray.) H 446 p 161

136. TEREBRA FLAVA, Gray.
Terebra flava, Gray, Proc. Zool. Soc. 1844, p. 60; Hinds, Thes. Conch. p. 177. no. 77, pl. 44. f. 75. DESH 1857 p 301
Hab. —?

80. *flava*. *Terebra* - GRAY, 1834, P. Z. S. L., p. 60. Hab. ?
Size: 1" (= 25.4 mm). Holotype: 23.4 mm (marked x inside aperture). = *Terebra lutescens* E. A. SMITH, C 69
1873.



Flava (*Terebra*), Gray. Proc. Zool. Soc., 1834, p. 60.
= *T. cancellata*, Quoy, 6/6

T. flava Gray = cancellata

TRUON 1885

6. *Terebra flava*, Gray (= *cancellata*). Conch. Icon., f. 126,

Terebra flava, Gray. p 203

The specimen from which the figure of this species in the 'Thesaurus Conchyl.' pl. xlv. fig. 75 was taken is in the Cumingian collection. On comparing it with the type of *flava* in the British Museum, it proves to be a very distinct shell; !! and I have therefore applied to it the name *lutescens*. SMITH 1873 p 203



Terebra flava Gray. P.Z.S. 1834, 60: Thes. Conch. I, 177, pl. 44, fig. 75.
Described without locality, the type being labelled "from an old collection". The holotype of *T. peasii* Deshayes, described from Hawaii in P.Z.S. 1859, 392 is identical with *flava*. TOMLIN 1851 40