

XIX: SYNONYME DE T. INCONSTANS HINDS

E. A. SMITH 1877
INCONSTANS

... sous-suturales. Blanc, tachée de pourpre sous la suture.

E 1294

Hastula inconstans Hinds, 1844. *H. confusa* (E. A. Smith, 1877) is a synonym.

ASN AUG 78

de *Lamarckia fusca* Martyn sous le nom de *cinerea*

2° Planche 121, figure du bas, à droite, constamment de plus petite taille, nettement et régulièrement plissée longitudinalement, dont l'ouverture, plus courte, s'élargit et est subtronquée à la base. Cette seconde espèce a été séparée par E. A. Smith, qui l'a nommée *T. confusa*, mais il est regrettable que cet auteur n'ait pas cité la figure de Martyn qui la représente et il nous paraît utile de combler cette lacune en donnant ici une liste de références qui s'y rapportent avec certitude.



Ph. Des 1935 p 42-43

Terebra (Hastula) confusa, sp. nov.

Terebra cinerea, Hinds (not of Born), Sowerby's Thesaurus Conchyliorum, vol. i. pl. 45. f. 130.

T. aciculina (part), Reeve (not of Lamarck), Conchologia Iconica, vol. xii. f. 121, d; varieties, f. 121, c & f.

In the 'Annals and Magazine of Natural History,' 1873, vol. xi. p. 262, I made a few remarks on a species of *Terebra* which had been referred by Messrs. Deshayes, Hinds, and Reeve (partly) to *T. aciculina*, Lamarck. Since writing the paper referred to I have had occasion to reexamine this species, and find that it is not in fact the Lamarckian shell. In his diagnosis, which appears almost as if it were extracted from Born's description of *T. cinerea* ('Index Musei Vindobonensis,' p. 267), from the similarity of language, he gives "albido-nereia" as the colour of the species, which evidently in no way applies to the present species, which is of a more or less hy-brown colour, banded with white beneath the suture, and so longitudinally streaked with brown, and has a white zone on the last whorl a little below the middle, which is distinctly seen within the brown aperture.

M. Deshayes, in the second edition of Lamarck's 'Animaux sans Vertèbres,' vol. x. p. 250, remarks that Lamarck was wrong in changing Born's name *cinerea* to that of *aciculina* if the shell was the same as that of this author, and he considers Born's species identical with that described by Lamarck under the name of *Terebra caeruleascens*. This, however, is a palpable mistake; for the latter is a smooth shell without any striation or plication beneath the suture, differently coloured, and has the whorls indistinct ("suturis obsoletis," Lamarck), which results from the fact of the whorls being encircled by a narrow white line just above, but contiguous to, the suture, which similarity induced me to found the subgenus *Impages* for this and a few other species possessing the same characteristics.

E. A. SMITH 1877 p 229

Confusa (*Terebra*), Smith. Ann. Mag. N. H., 1877, p. 229, pl. xix. = *T. aciculina*, Reeve, ... 32

15. *Hastula inconstans confusa* (Smith) ...

Height 35 mm.; diam. maj. 9 mm., min. 8 mm.




PLATE 8, BOTTOM LEFT

*Hastula confusa*³ (E. A. Smith), 1877
Length: 33 mm.

Distribution: Hawaii and Tropical Pacific
Habitat: Shallow water (common)

- 3. Shell shiny; longitudinally, straightly plicate, the plicae regular, sharp, extending across the whorls; aperture broadly dilated at the base; color variable from chocolate to off white, a white band anterior to the sutures and at periphery of the body whorl, the interstices of the ribs reddish chestnut on white shells; sometimes incorrectly called *inconstans* Hinds which is a synonym of *Terebra anomala* Gray.
- 1860. *Terebra aciculina* Reeve (not Lamarck), Conch. Ic., vol. 12, pl. 23, fig. 121d, var. c. and f.
- 1877. *Terebra confusa* E. A. Smith, Ann. Mag. N.H., p. 229.

Terebra confusa, E. A. Smith (= *T. aciculina*, Reeve, pars, Specimens from Singapore are in the British Museum (History).

TEREBRA CONFUSA E. A. Smith.

T. (Hastula) confusa E. A. SMITH, Ann. a. Mag. N. H., sér. XIX, p. 229.

aciculina Reeve, TRYON (pars, non Lamarck), Man., VII, p. 32 (non figuré).

— Iles Marquises (collect. Ph. D., ex D'Arbuthnot); (Papeete) Tahiti (collect. Ph. D., ex 890).

Des 1935 p 42

15. *Hastula confusa* (Smith). クチビロタケ H. 35, Dj. 9, Dn. 8.

HR STAKI 1112 Loc. Ogasawara.



WEAV 1960 vol 2408

SMITH . D 80 mm. Nord et ouest du Pacifique. Lignes de croissance, faibles côtes sous-suturales. Blanc, pourpre ou mélange des 2; sur les spécimens pourpres, petite zone blanche tachée de pourpre sous la suture.

E p 294

Hastula inconstans Hinds. 1844. *H. confusa* (E. A. Smith. 1877) is a synonym.

ASN AUG 78

de Lamarck fusca Martyn unis. cinerea

2° Planche 121, figure du bas, à droite, constamment de plus petite taille, nettement et régulièrement plissée longitudinalement, dont l'ouverture, plus courte, s'élargit et est subtronquée à la base. Cette seconde espèce a été séparée par E. A. Smith, qui l'a nommée *T. confusa*, mais il est regrettable que cet auteur n'ait pas cité la figure de Martyn qui la représente et il nous paraît utile de combler cette lacune en donnant ici une liste de références qui s'y rapportent avec certitude.

Terebra (Hastula) confusa, sp. nov. p 229

Terebra cinerea, Hinds (not of Born), Sowerby's Thesaurus Conchylorum, vol. i. pl. 45. f. 130.

T. aciculina (part), Reeve (not of Lamarck), Conchologia Iconica, vol. xii. f. 121, d; varieties, f. 121, c & f.

In the 'Annals and Magazine of Natural History,' 1873, vol. xi. p. 262, I made a few remarks on a species of *Terebra* which had been referred by Messrs. Deshayes, Hinds, and Reeve (partly) to *T. aciculina*, Lamarck. Since writing the paper referred to I have had occasion to reexamine this species, and find that it is not in fact the Lamarckian shell. In his diagnosis, which appears almost as if it were extracted from Born's description of *T. cinerea* ('Index Musei Vindobonensis,' p. 267), from the similarity of language, he gives "albido-cinerea" as the colour of the species, which evidently in no way applies to the present species, which is of a more or less ashy-brown colour, banded with white beneath the suture, and also longitudinally streaked with brown, and has a white zone around the last whorl a little below the middle, which is also distinctly seen within the brown aperture.

M. Deshayes, in the second edition of Lamarck's 'Animaux sans Vertèbres,' vol. x. p. 250, remarks that Lamarck was wrong in changing Born's name *cinerea* to that of *aciculina* if his shell was the same as that of this author, and he considers Born's species identical with that described by Lamarck under the name of *Terebra caerulea*. This, however, is a palpable mistake; for the latter is a smooth shell without any striation or plication beneath the suture, differently coloured, and has the sutures indistinct ("suturis obsoletis," Lamarck), which arises from the fact of the whorls being encircled by a narrow callosity just above, but contiguous to, the suture, which peculiarity induced me to found the subgenus *Impages* for this and a few other species possessing the same characteristic.

EA SMITH 1877 p 229

4. *Terebra confusa*, E. A. Smith (= *T. aciculina*, Reeve, pars, Lam.). Specimens from Singapore are in the British Museum (Natural History).

REVUE DES SCIENCES 1898 p 42 AMBROSIUM

Terebra confusa E. A. Smith.

Terebra (Hastula) confusa E. A. SMITH, Ann. a. Mag. N. H., sér. XIX, p. 229.

— *aciculina* Reeve, TRYON (pars, non Lamarck), Man., VII, p. 32 (non figuré).

localités. — Iles Marquises (collect. Ph. D., ex D'iel); Tahiti!; (Papeete) Tahiti (collect. Ph. D., ex iéret, 1890).

D2B & B6 1933 p 04

15. *Hastula confusa* (Smith). クチビロダケ H. 35, Dj. 9, Dn. 8

HR & TAKI 1112 Loc. Ogasawara

XIS: SYNONYME DE T. INCONSTANS HINDS 1844

EA SMITH 1877 228

INCONSTANS



AUBRY P 27

Ph. D2B 1935 p 42-43

Confusa (*Terebra*), Smith. Ann. Mag. N. H., 1877, p. 229, xix. = *T. aciculina*, Reeve, 32

TRYON 1881

WILSON 34

15. *Hastula inconstans confusa* (Smith)

Height 35 mm.; diam. maj. 9 mm., min. 8 mm.




PLATE 8, BOTTOM LEFT

*Hastula confusa*³ (E. A. Smith), 1877

Length: 33 mm.

Distribution: Hawaii and Tropical Pacific
Habitat: Shallow water (common)

3. Shell shiny; longitudinally, straightly plicate, the plicae regular, sharp, extending across the whorls; aperture broadly dilated at the base; color variable from chocolate to off white, a white band anterior to the sutures and at periphery of the body whorl, the interstices of the ribs reddish chestnut on white shells; sometimes incorrectly called *inconstans* Hinds which is a synonym of *Terebra anomala* Gray.

1860. *Terebra aciculina* Reeve (not Lamarck), Con. Ic., vol. 12, pl. 23, fig. 121d, var. c. and f.

1877. *Terebra confusa* E. A. Smith, Ann. Mag. N.H., p. 229.

WEAV 1900 Vol 240 8

