




TEREBRA MICANS. *Ter. testâ conico-subulatâ, acuminatâ, semiopacâ, pallidè fulvâ, nitidâ; anfractibus planulatis, integris, longitrorsum plicis capillaribus, supernè cæruleo et ad basin anfractûs ultimi fusco fasciatis; aperturâ infernè effusâ; columellâ truncatâ.*
 Axis 13 lin.
 Hab. — ?
 Cab. Cuming.
 The specific name I find in use as a cabinet name, but I am ignorant who is the originator.

MICANS (1)
 (HINDS) 1844
 236
 = ACICULINA



91. **T. MICANS.** (pl. xlv. f. 125.) *Hinds*, l. c. p. 157.

Testâ conico-subulatâ, acuminatâ, semiopacâ, pallidè fulvâ, nitidâ; anfractibus planulatis, integris, longitrorsum plicis capillaribus, supernè cæruleo et ad basin anfractûs ultimi fusco fasciatis; aperturâ infernè effusâ; columella nudâ, truncatâ.

Shell conico-subulate, acuminate, of a shining amber colour; whorls entire, polished, clothed with numerous very fine capillary folds, somewhat disappearing towards the lower part of the whorl; variously coloured with reddish-black, generally with a narrow band near the suture, and with the inferior whorls more coloured than the upper.

Native country unknown.

H. 1844b. T.E. p. 181 t. xlv fig. 125.

VOIR ACICULINA (BOUCHET 1982 p. 104).



HASTULA MICANS (Hinds, 1844) 25 mm - Atlantico-Occidentale - Sénégal - Cameroun - Côte d'Ivoire - In aqua bas...

AUSEY PL 3

Pale buff to grayish tan, with or without purplish-brown markings; to approx. 32 mm.

Hastula micans (Hinds)
 WEST AFRICA

Terebra micans (HINDS).

1910. *Terebra micans* Hinds, DAUTZENBERG, Contrib. F. Afr. Occid., I, p. 15.

Campagne de 1909-1910. Banc de Crawford et île Roume (îles de Los); dragage lagune Ebrié; dragage Grand-Bassam; Grand Popo, Dahomey, plage; dragage devant Cotonou, 20 à 25 mètres; plage de Saint-Paul-de-Loanda.

D 28 1913 p 6

78. **TEREBRA MICANS**, Hinds.
Terebra micans, Hinds, Thes. Conch. p. 181. no. 91, pl. 45. f. 125.
 Hab. — ?.

E 62 6581 H 530
 DESH 1859 p 299

137. *micans*, *Terebra* - HINDS, 1844, P. Z. S. L., p. 157.
 Hab. ? ("Cape Palmas, West Africa" on label). Coll. Cuming. Size: 13 lin. (= 29.3 mm). Holotype no. 1968248: 30.4 mm; syntypes: 23.1 mm, and 21.8 mm.

C 63

Terebra micans Hinds.

= ACICULINA LAM 1822 (BOU 83)

- 757. *Terebra Arvan* Adanson, Voyage au Sénégal, p. 33, pl. IV, fig. 1.
- 843. — *micans* Hinds, Proc. Zool. Soc. of London, p. 181.
- 360. *Terebra* — Hinds, Reeve, Conch. Icon., pl. IX, fig. 35. → 135!

Campagne de 1907. Entre Dakar et Rufisque, 16 mètres; baie de Hann, 9-12 mètres.

Campagne de 1908. N'Diago; Salines de Moudjeran; Baie de Hann, à basse mer.

Dispersion. — Sénégal.

Cette espèce a été réunie à tort par Tryon au *T. cinerea* orn.

D 28 1910 p 61

272



5 mm

272. **Terebra micans** Hinds (= *l'Arvan* Adanson). Coquille de 25 à 30 mm., très brillante, ornée seulement de très fines rides transversales, bien visibles sur la moitié inférieure des tours. Coloration blanche, unie ou marquée de bandes spirales brunes et fauves. Connue de la Mauritanie à l'Angola, cette espèce peut souvent s'observer vivante, par centaines, sur le sable de la côte atlantique des environs de Dakar, à marée descendante.

NICHOLS 1970 p 133

8. **TEREBRA MICANS** Hinds. Mauritanie-Angola. Avg. 25-35mm. C. C 65 81



8

MICANS (2)

HINDS N44

236

= ACICULINA

Terebra micans Hinds.

- 157 *Terebra Arvan* Adanson p. 53, pl. 4, fig. 4.
- 343 *Terebra micans* Hinds p. 157.
- 359 *Terebra micans* Deshayes p. 291.
- 359 *Terebra Adasoni* Deshayes p. 291.
- 360 *Terebra micans* Reeve pl. 9, fig. 35.
- 377 *Terebra micans* Marrat p. 239.
- 390 *Terebra micans* Dautzenberg p. 164.
- 391 *Terebra micans* Dautzenberg p. 12.
- 910 *Terebra micans* Dautzenberg p. 15.
- 912 *Terebra micans* Dautzenberg p. 6.
- 927 *Terebra micans* Lamy p. 385.
- 942 *Terebra micans* Fischer-Piette p. 163, pl. 3, fig. 1.
- 947 *Terebra micans* Nicklés p. 14.
- 950 *Terebra micans* Nicklés p. 133, fig. 272.

Material:

Lagos, Nigeria. Harry Madsen leg. 3 spec.
Guinea. 2 shells.

Remarks: The largest specimen had the top of the shell broken off and measured 23.7 x 6.2.

Distribution: Mauritania to Angola. **KNUDSEN 1952 p 181**




Micans (*Terebra*), Hinds. Proc. Zool. Soc., 1843, p. 181.

= *T. cinerea*, Born, 9/64. 31

64. *Terebra micans*, Hinds (= *cinerea*). Thes. Conch. f. 125,

TRYON 1885



Species 135. (Fig. a, b, c, d, Mus. Cuming.)

TEREBRA MICANS. *Ter. testâ aciculari-subulatâ, nitente- albâ, interdum lutescente, purpureo-brunneo-fasciatâ, an- fractibus declivi-planatis, indivisis, supernè densè elon- gatè plicato-crenulatis; aperturâ parvâ, columellâ ap- pressè arcuatâ, brevi.*

THE GLITTERING TEREBRA. Shell acicularly subulate, shining-white, sometimes yellowish, banded with rich- brown, whorls slopingly flattened, undivided, densely elongately plicately crenuled round the upper part; aperture small, columella appressly arched, short.

HINDS, Pro. Zool. Soc. 1843, p. 181.

Terebra Adasoni, Deshayes.

Hab. Senegal.

The typical peculiarity of this and the preceding species consists in the whorls not being divided by any spiral grooves, but densely puckered from the suture with fine elongated crenules.

RV 1860