

TEREBRA NIMBOSA. *Ter. testâ elongatè conico-subulatâ, acuminatâ, lacteâ, strigis longitudinalibus nubeculatâ; anfractibus planulatis, lævigatis, politis, integris, infernè propè suturam albâ, angustè fasciatâ, ultimo fasciato; columellâ lævi, truncatâ.* Axis 25 lin.

Hab. —? Cab. Cuming.

H 442 p 151

SEP D2B 1935 AT HECTICA p 2

26. T. NIMBOSA. (pl. xlii. f. 21. (*Hinds*, l. c. p. 151.

Testâ elongatè conico-subulatâ, acuminatâ, lacteâ, strigis longitudinalibus nubeculatâ; anfractibus planulatis, lævigatis, politis, integris, infernè propè suturam albo angustè fasciatis, ultimo fasciato; aperturâ effusâ; columellâ lævi, truncatâ.

Shell elongated, conico-subulate, acuminate, milky white, figured with aggregated slightly waved pale brown strigæ, ceasing suddenly a little above the suture, thus appearing narrowly banded with white; whorls smooth, entire, the last elongated. Aperture oval, effuse, without the least appearance of a canal. Columella, smooth, straight, and truncate.

Habitat unknown.



66
T. nimbosa Hinds
= caerulea var

H. 442 T. C. p 159 t xlii f 21

71. TEREBRA NIMBOSA, Hinds.

Terebra nimbosa, Hinds, *Thes. Conch.* p. 159. no. 26, pl. 42. f. 21.

Hab. Nouvelle Hollande.

DZS H 1859 p 289

66. *Terebra nimbosa*, Hinds (= *caerulea*, var.). *Conch. Icon.*, f. 37.

Nimbosa (*Terebra*), Hinds. *Proc. Zool. Soc.*, 1843, p. 151.
= *T. caerulea*, Lam., var. . . . 31

TRAYON 1875

143. *nimbosa*, *Terebra* - HINDS, 1844, P. Z. S. L., p. 151.

Hab. ? Coll. Cuming. Size: 25 lin. (= 56.3 mm).

Lectotype: 58.8 mm; syntypes: 64.9 mm, and 58.7 mm (worn and faded), and 52.6 mm. = *Hastula*

hectica (LINNAEUS, 1758).

C69

~~Red~~ *nimbosa*, *Hinds* . . . | Two specimens.

| Red Sea.

COCKE 1885 p 325

Species 37. (Mus. Cuming.)

TEREBRA NIMBOSA. *Ter. testâ subulatâ, ad basin truncato-ventricosâ, griseo-abbâ, cinereo-fusco longitudinaliter nebuloso-undatâ, anfractibus declivi-planatis, lævibus; aperturâ parvâ, columellâ rectâ, truncatâ.*

THE RAINY TEREBRA. Shell subulate, truncately ventricose at the base, greyish-white, longitudinally nebulously waved with ash-brown, whorls slopingly flattened, smooth; aperture small, columella straight, truncated.

HINDS, *Proc. Zool. Soc.* 1843, p. 151.

Hab. Red Sea; M. Botta.

A grey-clouded species, of very simple character, encircled with a faint depression in the place so frequently occupied by a groove.

RV 1860



37.