


148. *parva*, *Terebra* (*Leiostoma*) BAIRD in BRENCHLEY, 1873, *Cruise H. M. S. Curaçao*, p. 435, pl. 37, figs. 5, 6. Hab. New Caledonia. Size: 8-11 lin. \times 2 lin. (= 18-24.8 mm \times 4.5 mm). Lectotype no. 56: 22.8 mm (marked with an x on the tablet near the apex of the shell); paralectotype no. 65: 18.4 mm. C69

PARVA
BAIRD IN BRENCHLEY
1873



74. *Terebra parva*, Baird (= *cinerea*). *Voy. Curaçao*, t. 37, f. 6,

Parva (*Terebra*), Baird. *Cruise Curaçao*, Brenchley, 435, pl. 37, f. 5-6. = *T. cinerea*, Born, 174. 31

TREYON 1885

209. *Hastula parva* (Baird, 1873)

(Pl. 53, figs. 209a,b)

1873 *Terebra* (*Leiostoma*) *parva* Baird in Brenchley, *Cruise H. M. S. Curacao*, p. 435, pl. 37, figs. 5, 6.

Description: Shell to 24 mm; color shiny opaque white to bluish gray with evenly spaced yellow-brown subsutural spots; outline of whorls flat, slightly turreted; protoconch of 1½ yellow-brown whorls; sculpture almost smooth except for very slight crenulations immediately below suture; aperture moderately elongate; columella straight.


Type locality: "New Caledonia."

Distribution: Tropical Indo-Pacific; intertidal to 7 m.

Type: Lectotype BM(NH) no. 1979153; 22.8 mm.

Discussion: *H. rufopunctata* (210) may be separated from this species by its stripe at the periphery of the body whorl, and its translucent early whorls of the teleoconch with a stripe visible within.

6-209 - p 174



209a,b, *H. parva* (Baird in Brenchley). 209a, Paralectotype British Museum (N.H.); 18.4 mm. 209b, Upper whorls of paralectotype.