

5. *Euterebra inconspicua* PRITCHARD & GATLIFF.
Victoria.

INCONSPICUA
PRITCHARD & GATLIFF

1902 (120)

Euterebra Cotton and Godfrey 1932 (*inconspicua* Pritchard and Gatliff)
1005. *inconspicua* Pritchard and Gatliff 1902 (*Terebra*) (707, pl. 34, f. 11) KERSHAW 1927


= TRISTIS

KERSHAW



H.A. 659


x13/10



Pale brownish;
irregularly,
weakly ribbed;
to approx. 14 mm.
(BMNH type col-
lection 99242)

Euterebra inconspicua
Pritchard & Gatliff)
VICTORIA

34



120j

120j, Holotype of *T. inconspicua* Pritchard & Gatliff, National Museum of Victoria No. F-541; 10.8 mm.

BRAT 87

Terebra inconspicua, sp. nov. (Pl. IX., Fig. 2). 125

Shell small, not solid, of $9\frac{1}{2}$ whorls, regularly and tapering. Apex blunt. Nucleus of $1\frac{1}{2}$ smooth whorls. Whorls telescopically arranged, slightly convex with numerous faint ribs crossing the whorls at almost regular intervals, the only other sculpture being faint transverse striae, visible under the lens. Suture well-defined, and on account of the overlapping of the whorl a slight shouldering is caused. Colour sordid white, irregularly, longitudinally streaked with brown.

Dimensions.—Length, 12 mm., greatest breadth, 3.5 mm.

Locality.—Dredged five to six fathoms off Rhyll, Phillip Island, Western Port (J. H. Gatliff).

Observations.—This shell may be readily distinguished, owing to its slight sculptural characters.

Type in Mr. Gatliff's collection. PRITCHARD & GATLIFF 1902 p181



2