

INFLEXA

PEASE 1869

95

= SWAINSONI

I discover that the synonymy of *T. Peasii*, Desh., as published by me, was incorrect. I was led into the error by following the name attached to the typical specimens forwarded me. That species is without doubt a synonym of *T. puncticulata*, Desh., and the species mistaken for it is a variety of *T. Swainsonii*, Desh., departing so widely from the type that I distinguish it by a distinct name, as follows:

*T. Swainsonii*, Desh., var. *inflexa*, Pease.

The ribs on the type are straight, and interstices, especially on the upper whorls, faintly and remotely striate transversely.

The ribs on the above variety are flexuous, interstices impressly striate or grooved throughout, and whorls shorter. The shell is usually longer. The figure and description of *T. Swainsonii* in Conch. Icon., are taken from a specimen of the above variety.

PEASE 1869 t. 64

Inflexa (*Terebra Swainsoni*, var.), Pease. A. J. C., v, pp. 64, 86, . . . . . 21

TEYON 1885

*Terebra Swainsoni*, var *inflexa* Pease

Amer Journ of Conch V 1869 p 64, 86. *T. Swainsoni* Reeve conch ic pl 22 fig 118.

95

Weniger schlank, grösser, blasser; gebogene Rippen; erinnert an *cerithina*, doch sind bei dieser die Rippen näher und schwinden auf den letzten Umgängen völlig.

Fundort: Sandwich Inseln.

PLAETENS & LANGENAUER 1911 h. 11