

LÖBBECKEANA
 DUNKER 1877
 (=LINA) (42)



42b, Holotype of *T. loebbeckeana* Dunker, Zoological Museum, Humboldt University Berlin; 75.0 mm.

27. *Terebra Loebbeckeana* Dkr.
 Testa gracilis, longa, alba, fascia pallide fusca lata interrupta cincta, anfractibus circa 30 sutura distincta divisio, parum convexis, oblique costulatis transversimque striatis, subgranosis, superne sulco divisio instructa; apertura angusta; columella parum contorta; canalis subcurvus. — Alt. 75, lat. vix 10 mill. DUNKER 1877 174



Terebra Loebbeckeana Dkr.
 Tab. nostra 5. fig. 17. 18.

Malak. Bl. vol. 24. pag. 74.
 Testa longa, gracilis, alba fascia lata interrupta pallide fusca cincta, anfractibus circa 30 sutura distincta divisio, parum convexis, oblique costulatis transversimque striatis, subgranosis, superne sulco divisio instructa; apertura angusta; columella parum contorta; canalis subcurvus. — Alt. 75 mill. diam. max. vix 10 mill.
 Unicum tantummodo specimen inter Mollusca Japonica exstat notis nonnullis Terebrae Limae Desh. respondens, sed altera ex parte ab ea satis discrepans. Testa enim differt primo habitu graciliore, secundo superficie minus aspera, tertio praesertim eo, quod costa spiralis infra suturam sulco perspicuo formata, non est granosa, sed plicata, ita ut plicae longitudinales sulco isto tantummodo interruptae appareant.

Haec species amico Theodoro Loebbecke Molluscorum sedulo scrutatori dicata est.
 DUNKER 1882 172

Loebbeckeana (*Terebra*), Dkr. Mal. Blatt., xxiv, p. 74, 1/50. 29



T. LÖBBECKEANA, Dunker. Pl. 8, fig. 50.
 Whitish, tinged with light yellowish brown, with a wide, faint band of the latter color; subgranose by rather faint intersecting striae; sutural band slightly separated by a groove.
 Length, 75 mill. TRAYNER 185 175 Japan.

T. loebbeckeana
 Dkr.

50. *Terebra Löbbeckiana*, Dunker. Moll. Mar. Japon., t. 5. f. 17.