

PACHYZONA

MÖRCH 1861 (138)

= FORMOSA



138d

138d,

Worn lectotype of *T. pachyzona* Mörch, University Zoological Museum, Copenhagen; 51.0 mm.

196. *Terebra pachyzona* Mörch. — *T. forma* *T. robustae*; anfractus zona tumida oblique costulata divisi; pars antica costata, angusta, immersa, unde anfractus ultimi medio canaliculati; costis postice nodiferis, medio coarctatis; anfr. ultimi laevigati, sculptura evanescente, maculis quadratis triseriatis, suturalibus parvis pallidioribus; anfr. ultimus basi obtuse angulatus; columella buplicata. Long. 72, lat. 15 mill. — Realejo spec. 3 trita. non M. 1861 Mal. Bl. p. 105

*Pachyzona* (*Terebra*), Mörch. Mal. Blatt., vii, p. 105, . 40

TRAYON 1885

*Terebra pachyzona* Mörch.

(Figure 37.)

138


*Terebra formosa* DESHAYES, 1857. Jour. Conchyl., vol. 6, p. 65, pl. 3, fig. 6. Panama.

*Terebra pachyzona* MÖRCH, 1860. Malak. Bl., vol. 7, no. 3, p. 105 (Aug.). Realejo Nicaragua.

TYPE MATERIAL. Three worn specimens. Length, 72 mm.; diameter, 15 mm.

REMARKS. All three specimens are in the Copenhagen Museum lot and are, as Mörch realized, worn shells. On the basis of the discussion by Campbell, 1963 (Veliger, vol. 5, no. 3, p. 102, pl. 12, figs. 5, 8-13; pl. 13, figs. 3-6), I conclude that Mörch's species is synonymous with that of Deshayes. Mörch perhaps was unaware of the latter form, as he makes no mention of *T. formosa* in his comparisons. *Terebra* (*Terebra*) *formosa* Deshayes, 1857.

KEEN 1966 Col. J. Ac. Sci. Oce. PAPUA N.G. p. 29



37 a

37