

TEREBRA PICTA. *Ter. testâ subcylindraceâ, turrîto-subulatâ, nitidâ, pallidè aurantiacâ, atro-fusco longitrorsum maculatâ vel nebulosâ; anfractibus rotundatis, supernè cingulo tuberculato, infrâ plico-costatis, interstitiis striatis; anfractu ultimo fasciato; aperturâ parvâ, atro-fuscâ; columellâ subrectâ.* Axis 15 lin.
Hab. San Nicholas, island of Zebu, Philippines.
Cab. Cuming.

H-442 p156

75. T. PICTA. (pl. xlv. f. 105.) *Hinds*, l. c. p. 156.

Testâ subcylindraceâ, turrîto-subulatâ, nitidâ, pallidè aurantiacâ, atro-fusco longitrorsum maculatâ vel nebulosâ; anfractibus rotundatis, supernè cingulo tuberculato, infrâ plico-costatis, interstitiis striatis; anfractu ultimo fasciato; aperturâ parvâ, atro-fuscâ; columellâ subrectâ.

Highly polished, with a pale brown ground, ornamented with longitudinal dark reddish-brown spots. The folds are short, and crowded with striated interstices. Aperture is small, but not quite formed; last whorl band with white.

San Nicholas, Island of Zebu, Philippines.

A 1344b.T.C p.176 & xlv f.105.




162. TEREBRA PICTA, *Hinds*.

Terebra picta, *Hinds*, *Thes. Conch.* p. 176. no. 75, pl. 45. f. 105.
Hab. Philippines.

DESH 1859 p 305

* (x) *T. (Decorihastula) picta* HINDS, 1844: *Proc. Zool. Soc. London for 1843*: p. 156; *Fig.*: HINDS in SOWERBY; *Thes. Conch.*, 1844, T., pl. 45, fig. 105. Very uncommon. Luzon; Alabat Island (Olivera; B no. 007).

BUREAU 64



ONLY BAPT 37

Species 112. (Fig. a, b, Mus. Cuming.)

TEREBRA PICTA. *Ter. testâ elongato-turrîtâ, subcylindraceâ, flavescente, cœruleo fasciatâ, purpureo-fusco longitudinaliter maculatâ aut nebulosâ, anfractibus convexis, sulco supernè divisâ, concentricè plicato-costatis, interstitiis transversè lineari-sulcatis; aperturâ parviusculâ, oblongâ, columellâ parum contortâ, carinatâ.*

THE PAINTED TEREBRA. Shell elongately turreted, somewhat cylindrical, yellowish, blue-banded, spotted or clouded longitudinally with purple-brown, whorls convex, divided by a groove at the upper part, concentrically plicately ribbed, interstices transversely linearly grooved; aperture rather small, oblong, columella but little twisted, keeled.

HINDS, *Pro. Zool. Soc.* 1843, p. 156.

Terebra cinctella, *Deshayes*.

Hab. Island of Zebu, Philippines (in sandy mud, at four fathoms); Cuming. Mouth of the Indus.

A prettily blue-banded purple-blotched shell, elongately turreted, but of the same cylindrical outline and sculpture as *T. larvaformis*.

RV 1860



8. *Terebra picta*, *Hinds* (= *undulata*). *Conch. Icon.*, f. 112 b.

T. picta Hinds undulata

TRAYON 1885

Picta (*Terebra*), *Hinds*. *Proc. Zool. Soc.*, 1843, p. 156. = *T. undulata*, *Gray*.

22

81. *Terebra picta* *Hinds*, 1844

(Pl. 22, figs. 81a, b)
(Color pl. D, fig. 22)

1844 *Terebra picta* *Hinds*, *Proc. Zool. Soc. Lond.* pt. 11:156; 1844 *Hinds* in *Sowerby, Thes. Conch.* 176, pl. 45, fig. 105; *Reeve, Conch Icon.* 12, pl. 22, sp. 112b only.

Description: Shell to 30 mm, slender; color shiny gray or tan blotched with brown; outline of whorls flat; protoconch of 2 whorls; subsutural band narrow, noded, convex, defined by shallow groove and deep round punctations between ribs; axial ribs sharp, arcuate, wide-spaced, beginning at nodes on band, about 19 to 23 on penultimate whorl; spiral grooves, about 5, in interspaces; body whorl with narrow white stripe at periphery; aperture elongate; columella straight.


Type locality: "San Nicholas, Cebu Is., Philippines."

Distribution: Philippines to W. Australia.

Type: Holotype BM(NH) no. 1968241; 26.6 mm (top whorls missing).

Discussion: The color of this species gives it a marbled look. It may be separated from *T. campbelli* (83) by its longer whorls and convex subsutural band defined by deep round punctations. *T. campbelli* is considerably more straight-sided.

B-21-190



81a, Holotype British Museum (N.H.) No. 1968241; 26.6 + mm. 81b, Middle whorls of holotype.