

156. *polygonia*, *Terebra* - REEVE, 1860, Conch. Icon., pl. 27, spec. 154. Hab. ? Coll. Taylor. Size of figure: 10.6 mm. Holotype no. 74.12.11.302: 10.7 mm. C69

T. polygonia REEVE. Pl. 3, fig. 28.3. Tamatave, dragage, 2 ex., max. long. 14,5 mm. Les deux ex. s'accordent en couleur et en sculpture à la figure donnée par REEVE (Conch. Icon. 12, fig. 154). La couleur est jaune-brune grisâtre avec un lustre soyeux, le sillon sutural est distinctement marqué comme une ligne claire interrompue de plis et entouré d'une zone indistinctement plus foncée, une bande subsuturale d'une couleur un peu plus claire que le ton fondamental; les plis sont 6-7 au dernier demi-tour. L'exemplaire de REEVE est d'un habitat inconnu, c'est donc un fait des plus intéressants que de savoir que cette espèce se trouve à Madagascar, d'autant plus, qu'elle ne semble pas avoir été signalée depuis qu'elle fut décrite par REEVE. ODHNER AM 19 AKK für 200p39



30

x1



x8

Very similar to *T. pellyi* (q.v.); however, the early whorls are not as angularly convex; to approx. 13 mm. (BMNH type collection)

Terebra polygonia Reeve
HABITAT UNKNOWN S.K.

Species 154. (Mus. Taylor.)

TEREBRA POLYGONIA. *Ter. testâ gracili-turritâ, subpyramidalî, fulvo-brunneâ, anfractibus declivi-planatis, sulco supernè divisîs, validè varicoso-costatis, costis subdistantibus, interstitiis concavis, lineari-sulcatis; aperturâ parvâ, columellâ ad basin productâ et paululum recurvâ.*

THE MANY-CORNERED TEREBRA. Shell slenderly turrited, somewhat pyramidal, fulvous-brown, whorls slopingly flattened, divided at the upper part by a groove, strongly varicosely ribbed, ribs rather distant, interstices concave, linearly grooved; aperture small, columella produced at the base, and a little recurved.

Hab. — ?

A minute fulvous-brown species, angled throughout with strong rather distant varicose ribs. RV 1860



154

109. *Terebra polygonia* Reeve, 1860
(Pl. 29, fig. 109)

1860 *Terebra polygonia* Reeve, Conch. Icon. 12: pl. 27, sp. 154.

Description: Shell to 11 mm; color light fulvous-brown with darker subsutural band and darker base of body whorl; outline of whorls convex, protoconch of 3½ slender whorls; subsutural band narrow, somewhat constricted, with weak nodes, band defined by groove; axial ribs rather sharp, wide spaced, 13 on penultimate whorl; spiral sculpture not evident; body whorl with groove at periphery; aperture moderately elongate; columella slightly recurved.

Type locality: None given.

Distribution: Unknown.

Type: Holotype BM(NH) no. 1874.12.11.302; 10.7 mm.

Discussion: The type locality and distribution are unknown, and no other specimen is known except the holotype. B-109-1108

109. *T. polygonia* Reeve. Holotype British Museum (N.H.) No. 1874.12.11.302.; 10.7 mm.



109

109. T.

15 *T. polygonia* RV

15. *Terebra polygonia*, Reeve. Conch. Icon., f. 154,

T. POLYGONIA, Reeve. Pl. 7, fig. 15.

Strongly varicosely ribbed, ribs rather distant, interstices concave, spirally linearly grooved, sutural band distinct; fulvous brown. Length, 10 mill.

Hab. unknown (Taylor Coll.).

I am not acquainted with this species. TAYLOR 85p13

Polygonia (*Terebra*), Reeve. Conch. Ic., 154, 1860, #15 . 23