

TEREBRA PULCHRA. *Ter. testâ turritâ, conico-subulatâ, acuminatâ, nitidâ, pallidâ; anfractibus subplanulatis, longitrorsum rectè plico-costatis, supernè lineâ impressâ cinctis, interstitiis lævigatis; anfractu ultimo pallidè lineato.* Axis 11 lin.

Hab. Marquesas; in seven fathoms.

Cab. Belcher.

Perhaps more nearly resembling *T. plicata* than any other species, from which, with a little care, the description will suffice to distinguish it. The specimens were collected at the Marquesas group of islands, which scarcely offer any particular novelty in any department of natural history; and the greatest exception will be found among *Terebra*, of which it has a few peculiar species, and also some interesting varieties of other well-known kinds. Indeed, though the group is by no means the metropolis of the genus, the species would seem to exist here under some peculiar circumstances. H-44a p151

81. T. PULCHRA. (pl. xlv. f. 129.) Hinds, l. c. p. 151.

Testâ turritâ, conico-subulatâ, acuminatâ, nitidâ, pallidâ; anfractibus subplanulatis, longitrorsum rectè plico-costatis, supernè lineâ impressâ cinctis, interstitiis lævigatis; anfractu ultimo pallidè lineato.

Perhaps more nearly resembling *T. plicata* than any other species, from which, with a little care, the description will suffice to distinguish it.

Marquesas; in seven fathoms.

H-1844b T.C. p. 178 ex xlv f 129

134. TEREBRA PULCHRA, Hinds.

Terebra pulchra, Hinds, *Thes. Conch.* p. 178. no. 81, pl. 45.

f. 129.

Hab. Iles Marquises.

D25A 1857 p309

161. *pulchra*, *Terebra* - HINDS, 1844, P. Z. S. L., p. 151.

Hab. Marquesas. Coll. Belcher. Size: 11 lin. (= 24.7 mm). Holotype: 24.7 mm. = *T. cerithina* LA-

MARCK, 1822 (juvenile specimen). C69

Terebra pulchra, Hinds.

The type of this species, presented to the British Museum by Sir E. Belcher, on comparison with *cerithina*, Lamk., proves to be but the early stage of that form. S717H1873 p263

TEREBRA PULCHRA Hinds.

1843. *Terebra pulchra*

HINDS, Proc. Z. S. L., p. 151.

1885. — *cerithina*

TRYON (*pars*, non Lamarck).

Man., VII, p. 12, pl. 2, fig. 14 (*tantum*).

Localité. — Iles Marquises, 7 brasses (Hinds).

D20 & 86 1933 p69

PULCHRA

HINDS 1844

= CERITHINA

Species 155. (Mus. Brit.)

TEREBRA PULCHRA. *Ter. testâ turritâ, conico-subulatâ, acuminatâ, nitidâ, pallidâ, anfractibus subplanulatis, longitrorsum rectè plicato-costatis, supernè lineâ impressâ divisâ, interstitiis lævigatis, anfractu ultimo pallidè lineato.*

THE BEAUTIFUL TEREBRA. Shell turreted, conically subulate, acuminated, shining, pale, whorls rather flattened, longitudinally straightly plicately ribbed, divided at the upper part by an impressed line, interstices smooth, last whorl faintly lineated.

HINDS, Proc. Zool. Soc. 1843, p. 151.

Hab. Marquesas (in seven fathoms); Hinds.

This shell, of which there is no other specimen than the original in the British Museum, I had overlooked until it was too late to figure it. It is a small, rather swollen shell, colourless, but characteristic. RV 1860



14

T. pulchra Hinds
= *cerithina*

14. *Terebra pulchra*, Hinds (= *cerithina*). *Thes. Conch.* f. 129.

Pulchra (*Terebra*), Hinds. Proc. Zool. Soc., 1843, p. 151.

= *T. cerithina*, Lam. p. 2 f. 14. 12

TR40W 1885