

17. TEREBRA PURA Desh. (Pl. V, fig. 8.)

122

T. testa elongato-subulata, angusta, candida, nitente, eburnea; anfractibus latis, vix convexiusculis, sulco impresso, obsoleto, divisis, primis tenuè plicatis, alteris lævigatis, substriatisve, ultimo elongato, attenuato; apertura elongato-angusta, antice late emarginata, columella subcylindracea, extus angulo crasso, decurrente soluta.

Hab. Zanzibar.

Par sa forme et ses caractères, cette espèce se rapproche du *Senegalensis*; elle est cependant proportionnellement plus étroite; elle est partout d'une blancheur d'ivoire, sans la moindre apparence de coloration. Les premiers tours sont finement et irrégulièrement plissés; ces plis s'effacent insensiblement et se réduisent, sur les derniers tours, en de simples accroissements un peu plus marqués. Le sillon transverse qui partage les tours en deux parties inégales, est peu profond et disparaît presque entièrement sur les derniers tours. Un caractère distinctif se montre aussi dans l'angle aigu et saillant qui circonscrit au dehors la base de la columelle. **DESH 1857 p 82**



8

32. TEREBRA PURA, Desh.

T. testa elongato-subulata, angusta, candida, nitente, eburnea; anfractibus latis, vix convexiusculis, sulco impresso, obsoleto, divisis, primis tenuè plicatis, alteris lævigatis vel substriatis, ultimo elongato, attenuato; apertura elongato-angusta, antice late emarginata, columella subcylindracea, extus angulo crasso, decurrente soluta.

Long. 69 mill., larg. 11.

Hab. Zanzibar.

Collection Cuming.

DESH 1859 p 281

163. *pura*, *Terebra* - DESHAYES, 1857, J. C. P., 6: 82, pl. 5, fig. 8. Hab. Zanzibar. Coll. Cuming. Size: Not given in 1857; (69×11 mm in 1859). Holotype: 64.7 mm (marked x inside aperture, also on whorl **CG9**)

32. *T. pura* Desh. *Ÿ. Conchyl.*, vi, 82, pl. v, f. 8. Zanzibar. Type B.M. **T 44**

above aperture and near apex of the shell on the tablet). Syntypes: 51.5 mm, and 45.8 mm. **CG9**

PURA
DESH 1857
= FATUA (199)



199b

199b. Lectotype of *T. pura* Deshayes. British Museum (N.H.) No. 197970; 64.7 mm. **BRAT 87**

Species 67. (Mus. Cuming.)

TEREBRA PURA. *Ter. testâ elongato-subulatâ, nitente-albâ, anfractibus latis, convexiusculis, supernè sulco impresso obsoleto divisis, obliquè concentricè plicatis, plicis subdistantibus, tenuibus, versus aperturam gradatim evanidis; aperturâ oblongo-ovatâ, columellâ subrectâ.*

THE PURE TEREBRA. Shell elongately subulate, shining white, whorls broad, rather convex, obsoletely divided at the upper part by an impressed groove, obliquely concentrically plaited, plaits rather distant, thin, gradually fading towards the aperture; aperture oblong-ovate, columella nearly straight.

DESHAYES, Journ. de Conch. 1857.

Hab. Zanzibar.

Of the same type as *T. Senegalensis*, from the opposite side of the African continent, the characteristics of the species being all very feebly developed. **RV 1860**



67.

Pura (*Terebra*), Desh. Jour. Conch., 1857, p. 82, pl. v, f. 8, ¹⁶/₁₀



30

T. pura
Desh

30. *Terebra pura*, Desh. Jour. de Conch., 2d ser., ii, t. 5, f. 8.

T. PURA, Deshayes. Pl. 12, fig. 30.

Whitish, shining; obsoletely banded at the suture by an impressed groove, obliquely longitudinally plaited, plaits rather distant, thin, gradually fading towards the aperture.

Length, 2.5 inches.

TRAYON 75 p 16 Zanzibar.

Terebra pura Desh

Deshayes Journ Conch VI 1857 p 82 pl 5 fig 8. Reeve conch ic fig 67

undort: Nach beiliegender Etiquette von den Cardagos-Islands,

lurch Robillard erhalten. Deshayes gibt Zanzibar als

Vaterland an.

MAATEWS & LAWKAVEL 1871 126