

ROSEATA (1)


ADAMS & REEVE

1850

(103)


= TEXTILIS

SEE RICHARDS AT  
TEXTILIS



x1

x4



TEREBRA ROSEATA  
(Adams & Reeve,  
1850) 25 mm. Sooloo  
Islands, Pacific  
Oceania. Length 10  
mm.

Aobey PCC

131. TEREBRA ROSEATA, A. Adams et Reeve.  
*Terebra roseata*, A. Adams et Reeve, Voy. du Samarang, p. 30,  
pl. 10. f. 24.  
Hab. Iles Sooloo.

DESH 1850 p 300

170. *roseata*, *Terebra* - ADAMS & REEVE, 1850, Voy. Sam-  
arang, Zool., p. 30, plt. 10, fig. 24. Hab. Sooloo Is-  
lands. Size of figure: 29.5 mm. Holotype: 29.2 mm;  
syntypes: 25.2 mm, 21.0 mm, and 6.8 mm  
(juvenile specimen).  
CG9

\*(x) *T. (Strioterebrum) roseata* ADAMS & REEVE, 1850:  
Voy. Samarang, Moll., p. 30, pl. 10, fig. 24.  
Rare. Bataan, Luzon (CM no. 127134). BURCH

*Terebra (Strioterebrum) roseata* Adams &  
Reeve, 1848

BUR 63 100

1850?!


Tan to pale mauve-brown; with about  
18-20 ribs; to approx. 30 mm. Left  
specimen is syntype of *exigua* Deshayes,  
a synonym; right, a syntype of *roseata*  
(BMNH type collection).

*Terebra roseata* Adams & Reeve

TROPICAL PACIFIC

S.K.

5. TEREBRA ROSEATA. Pl. X. Fig. 24. Ter. testâ subulatâ, anfractibus plicato-costatis, costarum interstitiis liris tenuibus clathratis, sulco prominulo infra suturas; pallidè roseâ.  
HAB. Sooloo Islands; from sandy mud, at a depth of about thirty fathoms.  
Of a uniform, delicate, transparent, pink colour.



A DASH & REEVE 1850 p 300

13. *Terebra (Strioterebrum) roseata* Adams and Reeve.

ADAMS and REEVE. Moll. Voy, Samarang. p. 30, Pl. 10, fig. 24.

REEVE. Conch. Ic. Vol. XII, *Terebra*, fig. 104.

TRYON. Man. of Conch. Vol. VII, p. 21, Pl. 5, fig. 90.

Stat. 104. Sulu harbour, Sulu-island. 14 M. Sand. 3 Spec.

This characteristic species has been collected at or near the original locality.

SCHMIDT 1913 p 370

Species 104. (Mus. Cuming.)

TEREBRA ROSEATA. *Ter. testâ obeso-aciculari, subcylindraceâ, eximie roseâ, anfractibus convexis, sulco pertuso supernè divisio, undique densè plicato-costatis, interstitiis lineari-sulcatis; aperturâ parvâ, suboblique contractâ, columellâ laminatâ carinatâ, contorto-recurvâ.*

THE ROSEATE TEREBRA. Shell stoutly acicular, somewhat cylindrical, delicate rose-colour, whorls convex, divided at the upper part by a pricked groove, closely plicately ribbed throughout, interstices linearly grooved, aperture small, somewhat obliquely contracted, columella laminated, keeled, twistedly recurved.

ADAMS and REEVE, Moll. Voy. Samarang, p. 30. pl. 10. f. 24.

HAB. Sooloo Islands; Belcher.

Elaborately sculptured and uniformly tinted throughout with a pure soft rose-colour.

RN 1860 104

Roseata (*Terebra*), Adams and Reeve. Moll. Voy. Sam., p. 30, pl. x, fig. 24, . . . . . 21

90. *Terebra roseata*, Ads. and Reeve. Reeve, Icon., f. 104.

*T. roseata*  
Abic. 1850

T. ROSEATA, Adams and Reeve. Pl. 5, fig. 90.

Interstices of the ribs linearly grooved, sutural band defined by a punctate groove; columella laminated, twisted; rose-color. Length, 28 mill.

+ 2400 85 624

Soolo and China Sea.



104

ROSEATA (2)  
ADAMS & REEVE 1850

(103) = TEXTILIS


**Terebra roseata** Adams & Reeve, 1850

p. 149 Plate 53 Figure 2

Shell up to 25mm in length, very similar to *T. polygyrata* Deshayes, but differs in being larger, broader and less turreted. In mature specimens of *T. roseata* the columella is biplicate and the columellar callus-edge is risen above the body whorl surface. The species lacks the white sutural bands of *T. polygyrata* and is uniformly brown with a distinct pale mauve tinge, axial ribs are pale lilac in colour, protoconch brown; faded specimens are rose-coloured.

Moderately rare. Tropical Indo-Pacific; subtidal. *T. exigua* Deshayes, 1859, is a synonym. c70 p. 149

2. *T. roseata* Adams & Reeve. Indo-Pacific. To 1 inch. P. 148



103a

103a, Questionable type of *T. roseata* Adams & Reeve, British Museum (N.H.) No. 1979131; 29.9 mm.

BRAT 87