

**TEREBRA RUDIS.** *Ter. testâ turrîtâ, subulatâ, longitudinaliter plicatâ, spiraliter sulcatâ, cancellatâ; anfractibus planis, cingulo posteriore convexiusculo, noduloso; aperturâ mediocri; pallidè flavâ, apice flavo.*

Axis  $1\frac{1}{2}$  unc.

G. 34. p. 60

43. **T. RUDIS.** (pl. xliii. f. 60.) *Gray*, l. c. p. 60. *T. Petitii*, *Kiener*, l. c. p. 37, t. 13, f. 32.

Testâ elongatè conico-subulatâ, acuminatâ, turrîtâ, pallidâ; anfractibus planulatis, subangustis, supernè sulco divisis; plicis subrectis, brevibus, lineis duabus vel tribus impressis decussatis; aperturâ utrinque attenuatâ; columellâ nudâ, contortâ; breviter canaliculatâ.

So closely allied to *T. plicata*, that description must fail to distinguish them; yet on comparison, they are conspicuously distinct. The present is more elongated and coarser, the whorls are narrow and provided with short solid ribs, nearly or quite straight; and these are crossed by two, or perhaps three transverse lines. *T. plicata* is highly polished, the ribs are longer, delicate and sharp, placed somewhat at a distance, and very regularly crossed by three or sometimes four lines.

New York and Philadelphia: *Kiener*.

H. 19446. T. E. p. 165 t. XLIII f. 60.



69

45. **TEREBRA RUDIS**, *Gray*, l. c. p. 60.

*Terebra Petitii*, *Kiener*, l. c. p. 37. t. 13. f. 32.

*Hab.* Shores of New York and Philadelphia: *Kiener*. (v. s. in m. *Gray*.)

H 4442 p 163

SEE: DISLOCATUS (DALL 1889 p. 64.)

143. **TEREBRA RUDIS**, *Gray*.

*Terebra rudis*, *Gray*, Proc. Zool. Soc. 1834, p. 60; *Hinds*, Thes. Conch. p. 165, no. 43, pl. 43. f. 60 (exclusa *T. petiti*, *Kiener*).

*Hab.* —?

DESA 1259 t. 302

171. *rudis*, *Terebra* — *GRAY*, 1834, P. Z. S. L., p. 60.

*Hab.* ? Coll. J. E. *Gray*. Size:  $1\frac{1}{2}$ " (= 38.1 mm).

Holotype no. 74.11.10.11: 36.8 mm. = *T. dislocata* *SAY*, 1822.

C 69



185b

185b, Holotype of *T. rudis* *Gray*, British Museum (N.H.) No. 1874.11.10.11.; 36.8 mm.

BRAT 87

RUDIS  
GRAY 1834  
185  
= DISLOCATA

Species 34. (Mus. Cuming.)

**TEREBRA RUDIS.** *Ter. testâ subulato-turrîtâ, albidâ aut rufescente, apice flavo, anfractibus planis, longitudinaliter plicatis, transversim lineari-sulcatis, cancellatis, supernè sulco divisis, margine concentricè nodulatis; aperturâ parviusculâ, columellâ contortâ.*

THE RUDE TEREBRA. Shell subulately turreted, whitish or reddish, apex yellow, whorls flat, longitudinally plicated, transversely linearly grooved, cancellated, divided round the upper part by a groove, margin concentrically noduled; aperture rather small, columella twisted.

*GRAY*, Proc. Zool. Soc. 1834, p. 60.

*Myurella rufo-cinerea*, *Carpenter*.


*Hab.* South Carolina; *Dr. Jay*.

It is somewhat doubtful whether this is not a variety of *Say's T. dislocata*, figured above it.

KV 1860



34.



69. *Terebra rudis*, *Gray* (= *dislocata*). *Sowb.*, Thes., f. 60,

*T. rudis*  
*Gray*  
= *dislocata*

*Rudis* (*Terebra*), *Gray*. Proc. Zool. Soc., 1834, p. 60.

= *T. dislocata*, *Sav.* . . . . . 18

Revised 1840 1885