
Gastropoda: Caenogastropoda: Conoidea: Terebridae

Terebra spirosulcata Simone & Costa, 1999

Range: 22.1°S to 22.1°S; 41.07°W
Depth: 10 to 60 m (live 19 m)
Maximum Reported Size: 29.4 mm

Synonymy

Terebra brasiliensis auct. non E. A. Smith, 1873
Terebra doellojuradoi auct. non Carcelles, 1953
Terebra spirosulcata Simone & Costa, 1999

Details

Terebra spirosulcata Simone & Costa in Simone, 1999b, p. 237-241, figs. 1e-f, 3h, 6h-i, 7e, 8d-e, 13c, 25a-e, 26a-e

[Basis of the current name]

Type Locality: Off Ilha do Pai, Rio de Janeiro, Brazil, 30-40 m

Range: 22.1°S to 22.1°S; 41.07°W

Depth: 10 to 60 m (live 19 m)

Maximum Reported Size: 29.4 mm

Distribution: Brazil: Rio de Janeiro, Sao Paulo, Santa Catarina

Terebra brasiliensis auct. non E. A. Smith, 1873,

[Not available]

References: Simone (1999) s

Comments: Misidentification by Bratcher & Cernohorsky (1987, fig. 186c-d).

Terebra doellojuradoi auct. non Carcelles, 1953,

[Not available]

References: Simone (1999) s

Comments: Misidentification by Matthews et al. 1975.

Partial List of References

Simone, L. R. L. 1999. Comparative morphology and systematics of Brazilian Terebridae (Mollusca, Gastropoda, Conoidea), with descriptions of three new species *Zoosystema* 21 199-248.

Key To Reference Codes