

SULCATA

PEASE 1869

(95)

= SWAINSONI

Terebra sulcata Pease (Amer. Journ. Conch. V, p. 67, from Oahu) appears to be merely a small form of *swainsoni*, with stronger spiral sculpture, approaching that of *T. rosacea* somewhat. The specimens seen are No. 49967 M. C. Z., agreeing exactly with 117036 P. A. N. S., from 6-8 fathoms, off Honolulu. D. B. Langford.

PILSBRY 1921. p301

T. SULCATA, Pease. p. 62

T. aciculare, nitida; anfr. subangustis, infra suturas sulco cingulatis, longitudinaliter plicato-costatis, interstitiis concavis, profunde sulcatis, infra suturas striatis; apertura oblonga angusta; columella laevis, callo marginata, ad labium juncto; cinereo grisea, infra suturas pallidior, interdum costis pallidis.

Long. 15. Diam. 3 mill.

Hab. Insl. Oahu.

Shell acicular, shining; whorls rather narrow, encircled below the suture by a groove, longitudinally plicately ribbed, interstices concave, deeply grooved transversely, below the suture striated transversely; aperture oblong, narrow; columella smooth, nearly straight, bordered by a raised callousity, connecting with the outer lip; ashy grey, paler below the sutures, ribs occasionally pale ash color.

The above may prove to be a variety of *T. Swainsonii*, still farther removed from the type than *inflexa*, Pease.

PEASE 1869 p. 67

Sulcata (Terebra), Pease. Am. Jour. Conch., v, p. 67.

? = T. Swainsoni, Desh., var. 21

TUFTON 1885