

Mitra (Terebrifusus) multiplicata Lea.


Pl. 5, f. 65 * reprod. de Lea.

1840. *Mitra multiplicata* H. LEA FOSS. Claiborne p. 101, pl. 4, f. 49. — *Terebra multiplicata* ALDRICH Prelim. Report. 446.*Testa fusiformis, potius angusta, axialiter costulata; costis interstitia aequantibus; ultimo anfractu oblongo; columella 14 plicis ornata.*

Cette espèce a beaucoup de ressemblance avec la *Terebra terebriformis* Conrad, de laquelle elle diffère par les plis columellaires. M. Lea la réfère au gen. *Terebra*, M. Conrad ne la cite pas. D E GREGORIO 1890 p 70

Loc. Claiborne.

Terebra multiplicata H. C. Lea
See *Terebrifusus multiplicatus* (H. C. Lea) PALMER 1932 1966 1942

*Terebrifusus multiplicata* (H. C. Lea)

p 308

Plate 53, figs. 3, 8, 11, 14; Plate 88, fig. 4
Terebra multiplicata H. C. Lea, 1841, Amer. Jour. Sci., vol. 40, p. 101, pl. 1, fig. 19; H. C. Lea, 1848, p. 106; Pace, 1902, Proc. Mal. Soc. Lond., vol. V, p. 110.

Mitra (Terebrifusus) multiplicata (Lea) de Gregorio, 1890, p. 77, pl. 5, fig. 65.

T. testâ sub-turritâ, elongatâ crassâ, transversè striatâ, longitudinaliter costatâ, costis maximis; anfractibus—, valde convexis; suturis impressis; basi striatâ; columella quatuordecim plicis minimis; aperturâ; ovatâ; canaliculo sub-recurvo.

Shell sub-turritid, elongated, thick, transversely striate, longitudinally costate, with very large costae; whorls —, very convex; sutures impressed; base striate; columella with fourteen very small folds; mouth ovate; channel small, somewhat recurved.

Length.—. Breadth .25 of an inch.

Remarks.—This species much resembles the *T. gracilis*, Lea, but may be easily distinguished from that shell by the folds on the columella, its larger size, and its more strongly defined ribs. The mouth is .25 of an inch long. Its apex seems to be very fragile, for, although I have several specimens, that figured is the most perfect.—[H. C. Lea, 1841].

This species has generally been regarded as *T. amamus* (Conrad). The form has been discussed under that species.

Dimensions.—Height, 12 mm.; greatest diameter, 6 mm., lectotype.

Lectotype.—No. 13160, Academy of Natural Sciences, Philadelphia, Pa.

Occurrence.—Lower Claiborne: locality 727.

PALMER 1932 COLL BRIDAL RE 7(31) p 308

- | | | |
|-----|--|-----|
| 3. | <i>Terebrifusus multiplicatus</i> (H. C. Lea) | 308 |
| | No. 3147, P. R. I. H., 6 mm.; g. d., 2.5 mm.
Lower Claiborne: loc. 727 | |
| 8. | <i>Terebrifusus multiplicatus</i> (H. C. Lea) | 308 |
| | Same specimen as fig. 3 | |
| 11. | <i>Terebrifusus multiplicatus</i> (Conrad) | 307 |
| | No. 3148, P. R. I. H., 13 mm.; g. d., 4.5 mm.
Lower Claiborne: loc. 727 | |
| 14. | <i>Terebrifusus multiplicatus</i> (H. C. Lea) | 308 |
| | Same specimen as fig. 11 | |