

*Terebra (Myurella) reticulata*, Sow.

1839. *Terebra reticulata* Sow., *Trans. Geol. Soc. London*, (2), V., Pl. XXVI, fig. 9.

1857. *Terebra lima* Desh., *Jour. de Conch* (2), II, p. 69, Pl. III, fig. 2.

1879. *Terebra indica* Martin, *Tertiärschichten auf Java*, p. 33, Pl. VI, fig. 12.

1915. *Terebra indica* Martin, *Tesch, Pal. von Timor*, V, p. 36, Pl. LXXVIII(6), fig. 76.

1925. *Terebra (Myurella) reticulata* Sow., *Vredenburg, Mem. Geol. Surv. Ind.*, L, pt. 1, p. 21.

*Material*.—Two specimens (K24/265).

*Distribution*.—Recent : Indo-Pacific Province (*T. lima*). Gaj of Kutch, Miocene of Java and Pliocene of Timor.

*Remarks*.—The ornamentation of this species is somewhat variable. Compared with the type of *Terebra (Myurella) indica* Martin, in the collection of the Geological and Mineralogical Museum at Leiden, the Quilon specimens do not show any distinction of specific importance, although *T. indica* has a slightly different ornamentation, consisting in the presence very delicate spiral lines on the posterior side of the sutural band. DEY 1961 792