

ABRETIA
 ADAMS H & A 1853

GENUS: AECOS

Sub-gen. ABRETIA, H. and A. Adams.

Whorls longitudinally ribbed; columella simple, straight, produced anteriorly; outer lip not sinuated at the fore part.

T.S. CERITHINA

perithina, Lam.
 fictilis, Hinds.
 lepida, Hinds.
 mera, Hinds.

nassoides, Reeve.
 pygmaea, Hinds.
 tenera, Hinds.

ADAMS H & A 1853
 p. 225

Sous-genre. ABRETIA, H. et A. Adams, 1853. — Tours à côtes longitudinales. Columelle simple, droite, prolongée antérieurement. Bord interne parfois réfléchi. Bord externe simple, non sinueux à la partie antérieure. Ex. : T. cerithina, Lamarck.

CHENU 1859

Section ABRETIA H. and A. Adams, 1853. (OF TEREBRA)

Shell small, sculpture uniform, but in the later whorls more or less obsolete, sulcus persistent, whorls axially evenly ribbed, not constricted; young acute, adult subcylindric; columellar lip callous: the posterior angle of the aperture compressed into a sort of channel. Type *Terebra cerithina* Lamarck. PAU 1908

Abretia H. and A. Adams, 1853, Gen. Rec. Moll. 1: 235. [non Rafinesque, 1815]. Type species, *Terebra cerithina* Lamarck, 1822, recent, Indo-Pacific. "Whorls longitudinally ribbed; columella simple, straight, produced anteriorly; outer lip not sinuated at the fore part. Examples: *cerithina* Lam., *fictilis* Hinds, *lepida* Hinds, *mera* Hinds, *nassoides* Hinds, *pygmaea* Hinds, *tenera* Hinds." B & C 1982