

APPROXIMATA

DESH 1859

N. 308

125. TEREBRA APPROXIMATA, Desh.

T. testa elongato-acuminata, turrita, angusta, longitudinaliter costata, costulis apice albis interstitiis flavicantibus; anfractibus numerosis planis, linea vix impressa, inaequaliter bipartitis; interstitiis costularum densissime transversim striatis; ultimo anfractu cylindraceo, basi depressiusculo, costulis evanescentibus; apertura elongato-subquadrata, intus albo-flavicante; columella arcuata, cylindracea, basi canaliculata, late profundeque emarginata.

Var. *Testa paulo angustiore, flavo-ferruginea.*

Long. 42 mill., larg. 8.

Terebra pertusa, var. c, Kiener, Icon. des Coq. Viv. pl. 11. f. 24 C.

Hab. —?

Ma Collection.

DESH 1859 p. 299

SEE REPRESENTATION OF HOLOTYPE IN

UNDULATA REEVE 1860a fig 83

125. *T. approximata*, Deshayes, variety of *T. undulata*, Gray. 1860

TYPE IN REEVE DES NINES ? (T44)

Terebra approximata Deshayes, 1859. No figure was published. Type locality unknown. Reeve considered this taxa to be a junior synonym of *T. undulata* Gray, 1834, a fairly common Indo-Pacific species. No specimens of this taxa were found in the Deshayes collection. BRANCH 77 VEL 91(2) p 39.



96
T. approximata Desh
= *cancellata* Gr.

SAME FIG-
AS REEVE.

Approximata (*Terebra*), Desh. Proc. Zool. Soc., 1859, p. 299. = *T. cancellata*, Quoy, 22

96. *Terebra approximata*, Desh. (= *cancellata*). Reeve, Conch. Icon., f. 83,

TR40N 1885

Terebra approximata Deshayes, 1859, Proc. Zool. Soc. Lond. pt. 27: 299. The type is lost, and the species was not figured. N. 303 BR 87