

TEREBRA BICINCTA. *Ter. testâ subulatâ, lævigatâ, nitidâ; anfractibus rotundatis, indivisis, longitrorsum plicatis, supernè areâ coarctatâ, transversim biseriatis super plicas minutè tuberculatis; plicis tenuibus, acutis, interstitiis lævigatis; anfractu ultimo concolore.*
Axis 12½ lin.

Hab. — ?

Cab. Cuming. Unique.

Remarkably and very distinctly characterized by the two rows of small tubercles which encircle the whorls. The shell is otherwise of an uniform white glassy colour, which might be attributable to its condition.

H. 1844 p. 150

N. DUB.

Terebra bicincta Hinds, 1844, Proc. Zool. Soc. Lond. 11: 150; 1844 Hinds in Sowerby, Thes. Conchyl. 1:175, pl. 44, fig. 72. The type is lost, and the locality is unknown. The figure somewhat resembles the Panamic *T. glauca* Hinds or the Indo-Pacific *T. pustulosa* E. A. Smith, except that both have brownish shells. The figure of *T. bicincta* is white, and Hinds stated that the artist added an extra row of pustules.

N. DUB.

67 81

71. *T. BICINCTA*. (pl. xlv. f. 72.) *Hinds*, l. c. p. 150.

Testâ subulatâ, lævigatâ, nitidâ; anfractibus rotundatis, lineâ impressâ divisâ, longitrorsum plicatis, supernè areâ coarctatâ, transversim biseriatis super plicas minutè tuberculatis; plicis tenuibus, acutis, interstitiis lævigatis; anfractu ultimo concolore.



Remarkably and very distinctly characterized by the two rows of small tubercles which encircle the whorls beneath the girdling. The shell is otherwise of an uniform white glassy colour, which might be attributable to its condition. Our skilful artist seems to have seen an additional series of tubercles; but except on the last whorl where there are three, I cannot discover more than two. The specimen is unique in the collection of Mr. Cuming. Native country unknown.

H 1844 B.T.C. p 175 t XLIV f 72.

105. *TEREBRA BICINCTA*, Hinds.

Terebra bicincta, Hinds, Thes. Conch. p. 175. no. 71, pl. 44. f. 72.

Hab. — ?

DESH 1857 p 295



T. bicincta
79
Hinds

79. *Terebra bicincta*. Hinds. Sowb., Thes., f. 72.

T. BICINCTA, Hinds. Pl. 5, fig. 79. (fig)

White; with thin longitudinal plicæ, and no spiral sculpture; sutural band tuberculate, and with the plicæ below it cut into a couple of revolving rows of tubercles. Length, 28 mill.

Habitat unknown.

Described from a unique specimen in the Cumingian Collection.

TAYLOR 85 p 20