

TEREBRA CANCELLATA. *Ter. testâ turrîtâ, subulatâ, sulcis spirâlibus frequentibus profundis, plicis longitudinalibus æqualibus subconfertis; anfractibus planis, sulco posteriore profundo; aperturâ subparvâ, labio interiore incrassato elevato; pallidè cinereâ.*

Axis  $1\frac{3}{4}$  unc.

G 34 p 62

105. T. CANCELLATA, Gray, l. c. p. 62.

"Testâ turrîtâ, subulatâ, sulcis spirâlibus frequentibus profundis, plicis longitudinalibus æqualibus subconfertis; anfractibus planis, sulco posteriore profundo; aperturâ subparvâ, labio interiore incrassato elevato; pallidè cinereâ."

This I have never seen, and I believe the original specimen to be lost. Its length is one inch and three-quarters. The name has been previously applied M. Quoy.

Native country unknown.

H-44-b T.C. p 185

85. TEREBRA CANCELLATA, Gray, l. c. p. 62.

*Hab.* —?

This species I have not seen. The name, being preoccupied, requires to be changed.

H-44-a p 165

TEREBRA CANCELLATA, Gray (nec Quoy et Gaimard), Proc. Zool. Soc. 1834, p. 61.

DESM 1839 p 317

Espèce douteuse non figurée et très-incomplètement décrite.

Cancellata (*Terebra*), Gray. Proc. Zool. Soc., 1834, p. 62, 39

TAYLOR 1885: UNREC.

CANCELLATA

GRAY 1834

N. DUB

N. DUB

*Terebra cancellata* Gray, 1834, Proc. Zool. Soc. Lond. pt. 2: 62. (non Quoy & Gaimard 1833): The type is lost, the species was not figured, and the name is a primary homonym.

N. DUB

52 87