

46. TEREBRA DISPAR, Desh.

*T. testa elongato-subulata, angusta, albo-flavidula, aliquantisper griseo-fasciata, longitudinaliter tenue plicata, transversim acutissime striata; anfractibus numerosis, planulatis, continuis, plicis in medio evanescentibus; apertura elongato-angusta; columella brevi, acuta, late profundeque basi emarginata.*

Long. 29 mill., larg. 6.

Hab. —?

Ma Collection.

DESH 1859 p 284

DISPAR

DESH 1859

11.006

N. DUB

TYPE IN ZOOLOGIE DES MINES ? (T.44)

\*(x) *H. (Hastula) dispar* (DESHAYES, 1859): Proc. Zool. Soc. London, p. 284, no. 46; Fig.: REEVE; Conch. Icon., 1860, T., pl. 25, no. 137.

This species is commonly collected at Broome, Western Australia (B no. 317), but has not been among Philippine specimens examined.

BUREH 64

SHOULD BE INTERESTING TO KNOW WHICH SHELL BUREH COLLECTED.

*Terebra dispar* Deshayes, 1859. Figured by Reeve, pl. 25, fig. 137. Type locality unknown. Reeve said, "shining-white, semi-transparent. Very closely resembles *T. micans*." Tryon said it equals both *cinerea* (Born, 1778) and *aciculina* (Lamarck), two species which are quite distinct from one another. No specimens found in the Deshayes collection.

BENTON 77 VOL. VIII (2) 640

N DUB

*Terebra dispar* Deshayes, 1859, Proc. Zool. Soc. Lond. pt. 27: 284. The type is lost and the species was not figured. Reeve (1860) stated that it is similar to *micans*, and Tryon thought it was a synonym *cinerea* Born. 82 17

Dispar (*Terebra*), Desh. Proc. Zool. Soc., 1859, p. 284.  
= *T. cinerea*, Born, . . . . . 31



*T. dispar* Desh.  
= *cinerea*.

TRYON 1885

70. *Terebra dispar*, Desh. (= *cinerea*). Reeve, Conch.

Species 137. (Mus. Cuming.)

TEREBRA DISPAR. *Ter. testá aciculari-subulatá, nitente-albá, semipellucidá, anfractibus declivi-planatis, indivisis, supernè densè plicato-crenulatis; aperturá oblongo-ovatá, columellá appressè arcuatá, breviusculá.*

THE UNLIKE TEREBCRA. Shell acicularly subulate, shining-white, semitransparent, whorls slopingly flattened, undivided, densely plicately crenuled round the upper part; aperture oblong-ovate, columella appressly arched, rather short.

DESHAYES, Proc. Zool. Soc. 1859, p. 284.

Hab. —?

This species bears very closely on *T. micans*. It is more slenderly convoluted, and the crenules are stronger and shorter.

RV 1860

