

28. TEREBRA FUSCO-MACULATA, Sowerby, Tank. Cat. p. 23.

Hab. Senegal: Sowerby.

I cannot identify this shell, but there is much probability that it may be one of the varieties of *T. senegalensis*, Lamarck; a shell not so well known twenty years since as at present. 444a p 62

FUSCOMACULATA

Sow 1825

N. DUB

N-DUB

TEREBRA FUSCOMACULATA, Sow. Tank. Cat. p. 23. DESH 1825 p 318
D'après M. Hinds ce serait un double emploi de la *T. senegalensis*.

1985. TEREBRA-FUSCO MACULATA: p.

T. testâ turrito-subulatâ, acutissimâ, pallidè fuscâ, anfractibus non ventricosis, lineâ impressâ supernè divisâ, superioribus obliquè longitudinaliter sulcatis, inferioribus læviusculis, fusco-maculatis; areâ inter suturam et lineam impressam sulcatâ, fusco-maculatâ; long. 4 unc. lat. $\frac{1}{2}\frac{3}{8}$ unc.

Obs. We have formerly received this shell from Senegal, wherefore we at first supposed it might be Lamarck's *T. Senegalensis*; upon examination, however, it does not accord with his description. Our shell is of a light brown colour, the upper part of the volutions above the impressed line are regularly spotted with dark brown; the spots on the lower part of the volutions are arranged in rows, the upper of which is contiguous to the impressed line. The grooves of the upper volutions are interrupted by the impressed line. In the larger specimens the grooves become obsolete on the lower volutions.

Sow 1825 App p xxiii

Fuscomaculata (Terebra), Sowb. Tank. Cat., p. 23.
= *T. Senegalensis*, Lam. 16

TR40N 1885

Terebra fuscomaculata Sowerby, 1825, Cat. Shells Coll. Tankerville, App. xxiii. The type is lost, and the species was never figured. The type-locality is unknown. 1884

N DUB