

RUSTICA  
HINDS 1844  
N. DUB

TEREBRA RUSTICA. *Ter. testâ obeso-subulatâ, acuminatâ, fuscâ, nitidâ, striis transversis scabrâ; anfractibus subrotundatis, longitrorsum plico-costatis, supernè infra suturam luteis; plicis subdistantibus, ferè continuis; anfractu ultimo elongato, concolore; aperturâ elongatâ; columellâ laevi, subrectâ. Axis 8 lin.*  
*Hab. — ?*  
Cab. Metcalfe.

H. 442 p 158

N. DUB  
EXCLUENDAE ??


*Terebra rustica* Hinds, 1844, Proc. Zool. Soc. Lond. 11: 158.  
This was described from the Metcalfe collection, the whereabouts of which are unknown. The type locality is unknown and the figure indeterminable. Tryon (1885) listed it as *T. fulgurata* Philippi, which has since been placed in the Columbelloidea, but the aperture appears to be too flaring for that genus. It somewhat resembles the form of *T. armillata* Hinds that Carpenter described as *T. albocincta*.  
N. DUB

98. **T. RUSTICA.** (pl. xlv. f. 113.) *Hinds*, l. c. p. 158.  
*Testâ obeso-subulatâ, acuminatâ, fuscâ, nitidâ, striis transversis scabrâ; anfractibus subrotundatis, longitrorsum plico-costatis, supernè infra suturam luteis; plicis subdistantibus, ferè continuis; anfractu ultimo elongato, concolore; aperturâ elongatâ; columellâ nudâ, laevi, subrectâ.*  
An acuminate, retuse species, with distant straight rounded folds, and scabrous from crossing striæ. The aperture and last whorl are both elongated, and the latter displays no band. The colour is of a rusty red. The figure represents a fine specimen in the collection of Wm. Metcalfe, Esq.  
Native country unknown.

H. 442 p. 158 t. xlv f. 113



167. **TEREBRA RUSTICA**, Hinds.  
*Terebra rustica*, Hinds, Thes. Conch. p. 183, no. 98, pl. 45, f. 113.  
*Hab. — ?*  
DUSH 1855 p 306



19, 20. *Terebra rustica*, Hinds (= *fulgurata*).  
Icon., f. 147 a, b.


Species 147. (Fig. a, Mus. Metcalfe, Fig. b, Mus. Cuming.)

TEREBRA RUSTICA. *Ter. testâ fusiformi-oblongâ, acuminatâ, aurantio-rufâ vel cinereâ, infra suturas albo et purpureo obscure fasciatâ, anfractibus declivi-convexis, indivisis, varicoso-costatis, costis subdistantibus, interstitiis lineari-sulcatis; aperturâ oblongâ, columellâ carinatâ, vix contortâ.*

THE RUSTIC TEREBRA. Shell fusiformly oblong, acuminate, orange-red or ash-colour, obscurely banded beneath the sutures with white and purple, whorls slopingly convex, undivided, varicosely ribbed, ribs rather distant, interstices linearly grooved; aperture oblong, columella keeled, scarcely twisted.

HINDS, Proc. Zool. Soc. 1843, p. 158.  
*Hab. — ?*

There is a very near relationship between this species and some abnormal varieties of *T. fulgurata* and *aciculata*. The basal columellar keel is often conspicuously coloured in all three.



Rustica (Terebra). Hinds. Proc. Zool. Soc., 1843, p. 158.  
= *T. fulgurata*, Phil. . . . . 11/19-20 . . . . . 38

TRYON 1885