

Buccinum acus, Gmel. p. 3502. no. 141. ¹⁷⁹¹

Acus sartoria, Martini, t. 4. f. 1451. Espèce indéterminable par la défectuosité de la figure; il n'est pas bien certain pour nous qu'elle appartienne au genre *Terebra*. ^{1853 p 316}

Acus. 141. *B. testa acuta exalbida lineis undulatis horizontalibus, anfractibus bifidis crenulatis rugosis, columella spiraliiter torta.*

Martini. Conch. 4. t. 154. f. 1449.

Habitat . . . *testa 1½ pollicem circiter longa.*

GML 1791 b 3502

ACUS

GML 1791

EXCL.

EXCEPVERDAE

BOSC 1801

EXCL.

Vis pointue, *Terebra acus*.

Aiguë, très-blanche avec des lignes ondulées, horizontales; les tours de spire partagés, crénelés et rugueux; la columelle contournée en spirale.

Martini, Conch. 4. tab. 154. fig. 1449.
On ignore son pays natal. Bosc 1801

There still remain to notice the following species, which were referred by the old authors to that section of *Buccinum* which is now regarded as synonymous with *Terebra*, but which there is much difficulty and doubt in referring to any shells at the present time in our collections:—*Buccinum murinum*, Linn.; *B. acus*, Chemnitz; *B. hecticum*, Gmelin; *B. bifasciatum*, Petiver; *B. concinnum*, Gualtier; *B. succinctum*, Chemnitz; *B. geminum*, Linn.; *B. proximum*, Linn.; *B. monile*, Linn.; *B. sinuatum*, Linn.; and *B. phallus*, Chemnitz, is since known as *Pleurotoma buccinoides*. Hn 3 442 p 168

Acus (Buccinum), Gmelin. Syst. Nat., p. 3502, No. 141, . 39

UNDETERMINED - TROYON 1825

Acus. 148. Shell turreted, subulate; whorls transversely divided, crenulated, and wrinkled; pillar twisted spirally.

Buccinum Acus. Gmelin, p. 3502. Schreibers Conch. i.

p. 171.

Buccinum, No. 171. Schroeter Einl. i. p. 401.

Martini, iv. p. 297. t. 154. f. 1449.

Inhabits —

Shell near an inch and a half long, and about one sixth as broad, whitish, with horizontal undulated lines.

DML W-1817 b 649