

1. COLUMBELLA (ATILIA) ATTENUATA, Angas.

1871. *Columbella (Mitrella) attenuata*, Angas, Proc. Zool. Soc. p. 14, Species 4, pl. 1, fig. 4. 691

1883. *Columbella (Atilia) attenuata*, Tryon, Manual of Conchology, Vol. v. p. 151, pl. 53, fig. 18.

1884. *Terebra Beddomei*, Petterd, Journal of Conchology, Vol. iv. No. 5, p. 142, No. 28. →

1885. *Terebra Beddomei*, Tryon, Man. Conch. Vol. vii. p. 40.

Hab.—Sow and Pigs Reef, Port Jackson, 4 fathoms, Green Point, 8 fathoms, sand and shells (*J. Brazier*, 1864); Middle Harbour, in shell sand and shingle, Maroubra and Long Bays, near Sydney (*A. U. Henn*); Brown's River, Tasmania (*C. E. Beddome and W. F. Petterd*).

This is another variable species sometimes of a shining brown with a pale band at the sutures; others are finely mottled with network. Having Mr. Petterd's types before me from Mr. Beddome, I find that *Columbella attenuata*, Angas, is identical with Petterd's *Terebra Beddomei*. BRAZIER 1864 p. 691

EXCLUENDAE

BEDDOMEI
PETTERD 18

Beddomei (*Terebra*), Petterd. Jour. of Conch., iv, 142, 1884, NO 40

TRYON 1885

28. *Terebra Beddomei* n. sp. Shell elongate, white, smooth, apex blunt, suture scarcely impressed, whorls 6-7, very slightly convex. Long. 6½ mill., lat. 2 mill. Habitat, Brown's River. A plain form without ornamentation. PETTERD 1884 p. 142

PYRENE BEDDOMEI Petterd.

Columbella attenuata Angas, Proc. Zool. Soc., 1871, p. 14, Pl. i., fig. 4; Not *Columbella attenuata* Beyrich, 1854, Pace, Proc. Malac. Soc., v., 1902, p. 57.

Terebra beddomei Petterd, Journ. of Conch., iv., 1884, p. 142.

In the British Museum are five *attenuata* from Sydney, presented by Angas in 1871, and, therefore, probably types, but not so marked. Pace shows that this name was previously used by Beyrich in 1854, so it becomes necessary to revive Petterd's name, *T. beddomei*, generally acknowledged as a synonym. HEDLEY 1913 p. 321

NOT IN BEDDOMEI

NOT IN BAATCHEER'S