

100. T. COSENTINI. (pl. xlv. f. 107.) Philippi, Enum. Moll. Sicil. p. 227, t. 11. t. 22. ? → See Desh.: #28??

Testâ elongatè ovatâ, acuminatâ, turrîtâ, cinereâ, politâ; anfractibus integris, supernè tuberculis acutis uniseriatim coronatis, nigricante pictis, ultimo propè basin fasciato; aperturâ utrinque attenuatâ; columellâ nudâ, rectâ; canali brevi vel subnullo.

Shell ovate, acuminate, polished, the whorls crenated by a row of sharp tubercles, and between them and the suture is a much smaller series, not at first very conspicuous. The character of the shell is rather aberrant, and may be regarded as significant of its extra-tropical habitat.

Naples: Philippi.

N. 446. T. e. p. 184 t. xlv f. 107

EXCLUENDAE


COSENTINI 1836  
PHILIPPI 1836 -  
EXCL  
= ACICULATA NOT

NOT A  
TEREBRIDAE  
SEE BOUCHET.



105. TEREBRA COSENTINI, Philippi, p. 227. t. 11. f. 29.

Hab. Naples: Philippi. (v. s. in m. Cuming.) 1444a, 146



AGBR 1984. SEE TEXT

Terebra cosentini Philippi, 1836, Enum. Moll. Sicilae 1:227, pl. 11, fig. 29. This is a species of Mazatlanina from the west and east coasts of America (see Bouchet and Gofas, 1983).  
EXCL Bouchet 77

2. TEREBRA COSENTINI, Phil.

Terebra cosentini, Philippi, Enum. Moll. Sicil. t. 1. p. 227. pl. 11. f. 29; Hinds, Thes. Conch. p. 184. pl. 45. f. 107.

Hab. Tarento (Philippi).

Après avoir donné ce nom à cette espèce dans le premier volume de ses Mollusques de Sicile, M. Philippi la réunit à la T. aciculata dans le second volume du même ouvrage. Nous aurions suivi cet exemple si déjà plusieurs fois on ne nous avait mentionné cette forme dans le Méditerranée. Avant de supprimer l'espèce il est convenable d'attendre de nouvelles observations. Desh 1859 p 245

Cosentini (Terebra), Phil. Enum. Moll. Sicil., i, p. 227, pl. 11, f. 29. = T. aciculata, Lam. 38

TR40185

Species 152. (Mus. Cuming.)

TEREBRA COSENTINI. Ter. testâ obeso-fusiforâ, acuminatâ, albâ, fulvo-fusco latifasciatâ, anfractibus supernè declivi-angulatis, ad angulum muricato-tuberculatis; aperturâ oblongâ, subamplâ, columellâ rectâ, carinatâ.

THE COSENZA TEREBRA. Shell stoutly fusiform, acuminate, white, broadly-banded with fulvous-brown, whorls slopingly angled round the upper part, thickly tubercled at the angle; aperture oblong, rather large, columella straight, keeled.

PHILIPPI, Enum. Moll. Sicil. vol. i. p. 227. pl. 11. f. 29. Terebra nodosoplicata, Dunker.

Hab. Naples, West Africa, West Indies.

The evidence of this species inhabiting the Bay of Naples appears to be of very doubtful character.

RV 1860




152b.



152a.



17



16

T. Cosentini sh = aciculata  
T. Cosentini sh = aciculata

16, 17. Terebra Cosentini, Phil. (= aciculata). Reeve, Icon., f. 152 a, b,

37 TR40185