

EXCLUDED

HARRISONI
T. WOODS 77

EXC

- 3. TEREBRA (EURYTA) HARRISONI, Tension-Woods. 1893
- 1863. *Euryta pulchella*, Ad. and Angas, Proc. Zool. Soc. p. 418, pl. 37, fig. 14 (non *Terebra pulchella*, Desh.).
- 1865. *Euryta pulchella*, Angas, Proc. Zool. Soc. "Molluscan Fauna of S.A." p. 169, No. 81.
- 1875. *Euryta Brazieri*, Angas, Proc. Zool. Soc. p. 390, Pl. XLV. fig. 5, 5a (non *Terebra Brazieri*, 1871).
- 1877. *Mangelia Harrisoni*, Tenison-Woods, Proc. Royal Soc. Victoria, p. 56.
- 1884. *Daphnella Harrisoni*, Tryon, Man. Conch. Vol. vi. p. 306.
- 1885. *Terebra (Euryta) Angasi*, Tryon, loc. cit. Vol. vii. p. 38, pl. 12, fig. 26.
- 1886. *Terebra (Euryta) Angasi*, Tryon; Tate in Southern Science Record, p. 4.
- 1887. *Pleurotoma (Cithara) Harrisoni*, Gatliff in Victorian Naturalist, "List of some of the Shells of the Marine Mollusca found on the Victorian Coast," Vol. iv. No. 4, p. 59.
- 1893. *Euryta Angasi*, Tryon; Adcock, "A Hand List of the Aquatic Mollusca of S.A." p. 5, No. 125.

Hab.—Rapid Bay, South Australia, deep water (*G. French Angas*); South Australia (*Professor Tate and Adcock*); off Port Jackson Heads, 25 fathoms, sand floor (*J. Brazier*, 1874); Clark Island, Victoria (*Tenison-Woods*); Victoria (*J. H. Gatliff*): Port Stephens, N.S.W., on the ocean beaches after heavy weather; and in beach shingle and sand at Edward's Beach, Middle Harbour.

This species was first named by Adams and Angas *Euryta pulchella*; the specific name being already in use in *Terebra* by Deshayes, Mr. Angas in 1875 redescribed it from specimens that I sent him, under the name of *E. Brazieri*, he having already in 1871 named a *Terebra Brazieri*. Tenison-Woods in 1877 called it *Mangelia Harrisoni*. Mr. Tryon in 1885 renamed it *Terebra (Euryta) Angasi*. Mr. J. H. Gatliff of Melbourne sent me specimens in 1885, to have the name confirmed; at the time I wrote him out the history of the species, but in his list of the Victorian Mollusca he places it in *Pleurotoma*; it should be placed in the subgenus *Euryta* of *Terebra*; it is named in the Melbourne Museum by Tenison-Woods *Mangelia Harrisoni*; I have compared it with specimens from New South Wales. Woods' specific name must stand for this species. BRAZIER 1875 p. 393

Genus *Raphitoma*, Bellardi, 1847.

- RAPHITOMA HARRISONI, T. Woods.
- 1863. *Euryta pulchella*, Adams and Angas (non *Terebra pulchella*, Deshayes). P.Z.S. Lond., p. 418, pl. 37. f. 14.
- 1865. *Euryta pulchella*, Angas. *Id.*, p. 169.
- 1875. *Euryta brazieri*, Angas (non *Terebra brazieri*, Angas). *Id.*, p. 390, pl. 45, f. 5, 5a.
- 1877. *Mangelia harrisoni*, T. Woods. P.R.S. Vic., p. 56.
- 1884. *Daphnella harrisoni*, Tryon. Man. Conch., vol. vi., p. 306.
- 1885. *Terebra (Euryta) angasi*, Tryon. *Id.*, vol. vii., p. 38, pl. 12, f. 26.
- 1886. *Terebra (Euryta) angasi*, Tate. Southern Science Record, p. 4.
- 1887. *Pleurotoma (Cithara) harrisoni*, Gatliff. V.N., vol. iv., p. 59.
- 1894. *Terebra (Euryta) harrisoni*, Brazier. P.L.S., N.S.W., vol. ix., p. 693.

Hab.—Western Port. PATCHARD & GATLIF 1900 p. 179. (VICTORIA) 5

NOT IN B & C 87

NOT IN PATCHARD 2